

Root-induced Changes in the Rhizosphere of Extreme High Yield Tropical Rice: 1. Soil Chemical Properties

Erry Purnomo^{1*}, Maman Turjaman², Anna Hairani³, Athaillah Mursyid⁴, Dodik Choiron¹, Raina Yulia¹ and Mitsuru Osaki⁵

Received 11 August 2009 / accepted 10 March 2010

ABSTRACT

Root-induced Changes in the Rhizosphere of Extreme High Yield Tropical Rice: 1. Soil Chemical Properties (E Purnomo, M Turjaman, A Hairani, A Mursyid, D Choiron, R Yulia and M Osaki): Padi Panjang cultivar is one of many local rice cultivars found in South Kalimantan that yields 8 Mg ha⁻¹ without fertilizer after last transplanting. The mechanisms involved in sustaining nutrient supply to sustain the extreme high yield are of interest. The following work aims to investigate the changes of soil chemical properties in rhizosphere of Padi Panjang cultivar. The Padi Panjang cultivar was grown in a rhizobox filled with soils from 3 different villages in Banjar Regency, South Kalimantan Province, namely, Kuin, Bunipah and Guntung Papuyu. The rice plant was grown for 5 weeks. At the end of the growing period, soil chemical properties such as pH, aluminum (Al), phosphorus (P), potassium (K⁺), ammonium (NH₄⁺), and nitrate (NO₃⁻) were measured. The results showed that Padi Panjang cultivar had the capability to change the soil chemical properties in the rhizosphere. The impact was more extent compared with IR64 cultivar. The changes were depended on soil character, especially, soil texture. The soil from Guntung Papuyu was the least affected by root. It was observed that Padi Panjang cultivar acidified more than IR64. A depletion zone of K⁺ and NH₄⁺ was found in the rhizosphere of both Padi Panjang and IR64 cultivars. The depletion zone of these ions could reach as far as 3 cm from the rhizosphere. For P, the depletion zone only occurred in the rhizosphere soil of IR64 cultivar. However, for Padi Panjang cultivar, the depletion zone of P did not exist. The Padi Panjang cultivar was able to maintain P concentration the same as or higher than control soil without plant. This is the first report showing that Padi Panjang cultivar can be considered as efficient lowland rice cultivar in absorbing not only P but also K in a P- and K-deficient-soil.

Keywords: Local rice, nitrogen, phosphorus, potassium, Padi Panjang, South Kalimantan, tidal swamp

INTRODUCTION

The local farmers in South Kalimantan, Indonesia used to grow local rice in tidal swamp area. They have more hundreds local cultivars grown by the local farmers. Some of them yield more than 3 Mg ha⁻¹ without fertilizer (Hasegawa *et al.* 2004). Our previous study (Purnomo *et al.* 2010) found that there was a local rice cultivar, called Padi Panjang, yielded 8 Mg ha⁻¹. This local rice cultivar originated from Kuin village, Banjar Regency, South Kalimantan. We

did not find any relationship between soil chemical properties measured with yield of Padi Panjang cultivar. This indicates there may be other mechanisms driven the nutrient supply to sustain such an extreme high yield, such as rhizosphere function.

Rhizosphere is defined as a zone of soil under direct influence of plant root. Some studies had shown rhizosphere soil differs from bulk soil (Eo and Nakamoto 2006). Previous studies had shown the effect of root in influencing the rhizosphere soil. Compared to bulk soil, rhizosphere soil had lower

¹Program Study of Natural Resource and Environmental Management, Lambung Mangkurat University, South Kalimantan; e-mail: erry_purnomo@telkom.net. *Present address: Silviculture Program Study, Faculty of Forestry, Lambung Mangkurat University.

²Research and Development, Forestry Department, Bogor.

³Research Institute of Swampy Crop, Banjarbaru

⁴Faculty of Agriculture, Lambung Mangkurat University, Banjarbaru Campus, South Kalimantan 70714, INDONESIA.

⁵Laboratory of Rhizosphere Environment Technology, Division of Bio-systems Sustainability, Graduate School of Agriculture, Kita-9, Nishi-9, Kitaku, Sapporo 060-8589, JAPAN. E-mail: mosaki@chem.agr.hokudai.ac.jp

J Trop Soils, Vol. 15, No. 2, 2010: 95-102

ISSN 0852-257X

pH (Grinsted *et al.* 1982; Begg *et al.* 1994) and nutrient concentrations, such as NH₄, Ca, Mg, and K (Yanai *et al.* 2003), and P (Li *et al.* 1991; Gahoonia *et al.* 1994; Luster *et al.* 2009).

This work focuses on investigating the function of rhizosphere of the Padi Panjang cultivar. According to Hinsinger *et al.* (2005) the rhizosphere differs from the bulk soil in a range of biochemical, chemical and physical processes that occur as a consequence of root growth, water and nutrient uptake, respiration and

rhizodeposition. Therefore, this research aims to study the effect of rhizosphere on changes of soil chemical properties.

MATERIALS AND METHODS

Soil

Soils used for the experiment were collected from the three villages, namely, Kuin, Bunipah and

Table 1. Selected soil properties.

No.	Soil Properties ^{*)}	Soil origin ^{**)}		
		Kuin (3°22'24''S; 114°32'19''E)	Bunipah (3° 27'52''S; 114 °32' 54''E)	Guntung Papuyu (3 °27'44'' S; 114 °36'50'' E)
1.	Particle size analysis (%) ¹			
	Sand	2.08	2.51	0.83
	Silt	61.20	73.82	57.85
	Clay	36.72	23.67	41.32
	Texture	Silty clay loam	Silty loam	Silty clay
2.	Organic C (%) ²	4.86 (high)	3.56 (high)	3.24 (high)
3.	Total N (%) ³	0.28 (moderate)	0.24 (moderate)	0.32 (moderate)
4.	C/N	17 (high)	14 (moderate)	10 (low)
5.	P _{Bray 1} (mg kg ⁻¹) ⁴	3.51 (very low)	3.10 (very low)	9.22 (very low)
6.	P ₂ O ₅ (mg kg ⁻¹) ⁵	497 (high)	444 (high)	289 (moderate)
7.	K ₂ O (mg kg ⁻¹) ⁶	876 (very high)	630 (very high)	357 (moderate)
8.	pH H ₂ O ⁷	4.02 (very acidic)	4.28 (very acidic)	4.18 (very acidic)
9.	Exch.-Ca (cmol(+) kg ⁻¹) ⁸	4.47 (low)	3.89 (low)	3.94 (low)
10.	Exch.-Mg (cmol(+) kg ⁻¹) ⁸	5.48 (high)	6.63 (high)	6.05 (high)
11.	Exch.-Na (cmol(+) kg ⁻¹) ⁸	0.04 (low)	1.34 (very high)	0.32 (moderate)
12.	Exch.-K (cmol(+) kg ⁻¹) ⁸	0.25 (low)	0.32 (moderate)	0.09 (very low)
13.	KTK (cmol(+) kg ⁻¹) ⁹	32.50 (high)	27.25 (high)	39.00 (high)
14.	Base saturation (%)	44 (moderate)	42 (moderate)	43 (moderate)
15.	EC (dS m ⁻¹) ¹⁰	0.20	0.10	0.02
16.	Al saturation (%) ¹¹	2.15 (very low)	2.38 (very low)	0.50 (very low)

Note: ^{*)} Procedure of measurements are described in ¹ Gee and Boulder (1986); McLean (1982); ²Yeomans and Bremner (1988); ³Bremner and Mulvaney (1982); ⁴John (1970); ⁵Olsen and Sommers (1982); ⁶Knudsen *et al.* (1982); ⁷McLean (1982); ⁸Thomas (1982); ⁹Rhoades (1982a); ¹⁰Rhoades (1982b); ¹¹Exchangeable Al, Dougan and Wilson (1974). ^{**)The values obtained were categorized as described in Djaenuddin *et al.* (1994).}




Figure 1. The layout of rhizobox.

Guntung Papuyu. The soils properties were shown in Table 1.

Rhizobox

The experiment was carried out in glass using a home-made rhizobox developed by (Wang *et al.* 2002). The rhizobox illustration can be seen in Figure 1. The nylon cloth used can not be penetrated by plant roots, however, mycorrhiza and water can be easily penetrate. The rice plants were grown in the middle segment of the rhizobox.

Treatments

The treatments of the experiment are shown in Table 2.

Rice Cultivation

Twenty rice seeds were sown at the middle segment of the rhizobox (see Figure 1) under saturated soil condition. At the 3rd day, seedlings were thinned to 10. The high plant density was deliberately done to create a rhizosphere soil and bigger impact of root on the soil. Seven days after emerging, the rhizobox was filled with deionized water to 1 cm depth and maintained till the end of growing period. To protect from pest attack the rhizoboxes were covered with a mosquito net. The plants were grown for 5 weeks.

Soil Sampling

One day before soil sampling, the watering was stopped to let the soil dried out. Each of soil segment

Tabel 2. Treatments for the rhizobox study.

Treatments ^{*)}	Notes		1	2	3
Soil origin:	Exchangeable Na	Clay content (%)			
• Kuin	low	37	√		
• Bunipah	Very high	24		√	
• Guntung Papuyu	moderate	41			√
Rice cultivars:					
• Padi Panjang	Extreme high yield local cultivar		√	√	√
• IR64	Improved cultivar, as a comparison		√	√	√

Note: ^{*)} each treatment was replicated 4 times.

from each rhizobox was excavated, put in a plastic bag and kept in a refrigerator (4°C) until used.

Soil Analysis

Soil analyses were conducted prior to planting and after the growing period. Soil analyses prior to planting were carried out for characterizing the soils used in the experiment. The soil properties used in this experiment are demonstrated in Table 1. At the end of growing period soil properties such as pH, Al, P, K, NH₄⁺-N and NO₃⁻-N were analysed. Soil was extracted using a 0.01 N CaCl₂ according to methods described in Houba *et al.* (1994). The use of 0.01 N CaCl₂ is believed not only maintains the soil solution ionic strength but also is cheaper and environmental friendly. The pH, Al and P were determined using methods described in McLean (1982), Dougan and Wilson (1974) and John (1970), respectively. While, the NH₄⁺ and NO₃⁻ concentration in the extract was measured colorimetrically using methods described in Kempers and Zweers (1986) and in Yang *et al.* (1998), respectively.

Data Analysis

Standard errors were shown to indicate data variation as affected by treatments set up.

RESULTS AND DISCUSSION

Soil Properties

The properties of soils are presented in Table 1. It is pointed out here that for growing improved rice cultivar there will be at least 2 problems faced. These are the low P availability and very low soil pH. In addition, unpredicted water level often submerges the newly planted seedlings. Nevertheless, the potential nutrients reserve such as N, P, and K are generally high.

It can also be observed that there were some differences among the soils. Firstly, the texture of soil from Guntung Papuyu was heavier than the other two soils. Next, the soil from Guntung Papuyu possessed highest C to N ratio, followed by soils from Bunipah and Kuin, respectively. Lastly, the Na⁺ and K⁺ concentrations and EC reading of soil from Bunipah




Figure 2. Change of soil pH (a) and soil Al concentration (b) at various distances from the rhizosphere. Bars indicate standard error of mean.

were higher than of soils from Kuin and Guntung Papuyu. It was noticed that Bunipah located near the river, therefore, it was highly affected saline water resulted from the tidal movement. These soil properties differences may contribute to the difference in magnitude affect of root on to the soil properties.

Change in Soil pH

Figure 2a shows the change of soil pH from various distances from the rhizosphere. Except soil from Guntung Papuyu, pH rhizosphere soil of Padi Panjang cultivar was lower than of IR64 cultivar. Even, in soil from Bunipah, root of Padi Panjang cultivar acidified the soil more than to the control soil without plant. The lower soil pH in the rhizosphere of Padi Panjang cultivar may be explained as follow. Purnomo *et al.* (2005) observed that compared to IR64 cultivar, Padi Panjang cultivar had a larger rooting system. In addition, our unpublished data showed that Padi Panjang cultivar had much higher biomass compared to IR64 cultivar. So, it may be plausible that the root of Padi Panjang

cultivar (1) releases more soluble organic acid, (2) has more oxidative condition in the rhizosphere results in the oxidation of potential acidic materials, namely, soluble iron Fe^{2+} , (Saleque and Kirk, 1995), mangan (Mn^{2+}) and/or sulphur (S^0), and (3) takes up more cations than anions, therefore, H^+ is released by the root for maintaining neutrality of electricity charge of soil interface (Grinsted *et al.* 1982; Begg *et al.* 1994).

Change in Soil Al Concentration

For all soils, the Al concentration in soil grown with plant was lower compare with control soil without plant (Figure 2b). It was mentioned earlier that the release of exudates from the root may chelate the Al in the soil solution. The release of exudates decreased the Al concentration as far as 5 cm from the rhizosphere.

Change in Soil P Concentration

It can be observed that P concentration in rhizosphere of Padi Panjang cultivar was higher than




Figure 3. Change of soil P (a) and soil K (b) concentrations at various distances from the rhizosphere. Bars indicate standard error of mean.

of IR64 cultivar grown in soils from Kuin and Bunipah (Figure 3a). For IR64 cultivar, a depletion zone was noticed, especially for soil for Kuin. The P concentration in the soil rhizosphere was lower than soil outside the rhizosphere. Some works have shown a similar phenomenon occurred with maize (Gahoonia *et al.* 1994), white clover (Li *et al.* 1991) and a model developed by Luster *et al.* (2009).

For Padi Panjang, however, no depletion zone was observed. The P concentration was higher than in the rhizosphere of IR64 cultivar. In Kuin's soil, the P concentration of the rhizosphere soil was the same as of the soil only. Even, in Bunipah's soil, the P concentration of the rhizosphere soil was higher than of the soil only. Using different plant *Brassica napus* (Grinsted *et al.* 1982) found an increase in P concentration in the soil rhizosphere as a result of pH decrease. They suggested that *Brassica napus* is efficient in absorbing P from P-deficient. It may reasonable to consider that Padi Panjang cultivar is efficient in absorbing P. It was also observed that the root of rice affected the P concentration as far as 3 cm.

Changes in Soil K⁺ Concentration

Figure 3b shows the concentration of K at various distances from the rhizosphere. Different from P, it was observed the patterns of K concentration were similar for both rice cultivars, being lower in the soil rhizosphere and increasing as moving away from the rhizosphere. As with P, a depletion zone was also observed for K. Using rice plant, Li *et al.* (2002) also found a similar result. The present works showed that the depletion zone could be noticed up to 3 cm. Our results demonstrated K concentration in soil grown with Padi Panjang cultivar higher than in soil grown with IR64. This may be associated with the lower pH as result of exudates release that may dissolve the reserved K. The effect of exudates in dissolving the reserve K can reach up to 5 cm from the rhizosphere.

Change in Soil NH₄⁺ Concentration

The patterns of NH₄⁺ change for all soils were similar for both rice cultivars (Figure 4a). As for P and K, a depletion zone was also noticed. Comparing




Figure 4. Changes in (a) soil N-NH₄⁺ and (b) soil N-NO₃⁻ concentrations at various distances from the rhizosphere. Bars indicate the standard error of mean.

with soil only, NH_4^+ concentration of soil with plant was lower. The NH_4^+ concentrations were similar for Padi Panjang and IR64 cultivars. It was also observed that limit of NH_4^+ could be absorbed that was approximately 50-55 mg $\text{NH}_4^+\text{-N kg}^{-1}$. It is suggested that only free soluble NH_4^+ could have been absorbed by the plant.

Changes in Soil NO_3^- Concentration

The change of NO_3^- was less extent compared to NH_4^+ . The change of NO_3^- at various distances from the rhizosphere was hardly notice (Figure 4b). It is common that in submerged soil nitrification process is inhibited (Purnomo 1996). The similarity of NO_3^- concentration in the planted soil and soil only suggested that NH_4^+ was the main source of N for the rice crops in the current study.

CONCLUSIONS

The Padi Panjang cultivar had the capability to change the soil chemical properties in the rhizosphere. The impact was more extent compared with IR64 cultivar. The changes were depended on soil character, especially, soil texture. Soil from Guntung Papuyu was the least affected by root. It was showed that Padi Panjang cultivar acidified more than IR64. A depletion zone of K and NH_4^+ was found in the rhizosphere of both Padi Panjang and IR64 cultivars. The depletion zone could reach as far as 3 cm from the rhizosphere. For P, the depletion zone only occurred in the rhizosphere soil of IR64 cultivar. However, for Padi Panjang cultivar, the depletion zone of P did not exist. The Padi Panjang cultivar was able to maintain P concentration the same as or higher than control soil without plant. It is the first report showed that Padi Panjang cultivar can be considered as efficient lowland rice in absorbing not only P but also K in a P- and K-deficient-soil.

ACKNOWLEDGEMENT

We thank the Directorate General of Higher Education, Indonesian Ministry of National Education for financing the work. The University of Hokkaido for supplying the nylon cloth and chemicals is greatly appreciated. We also acknowledge the construction criticisms of Journal of Tropical Soils reviewers.

REFERENCES

- Begg CBM, GJD Kirk, AF Mackenzie and HU Neue. 1994. Root-induced iron oxidation and pH changes in the lowland rice rhizosphere. *New Phytol* 128: 469-477.
- Bremner JM and CS Mulvaney. 1982. Nitrogen-Total. In: AL Page, RH Miller and DR Keeney (eds), *Methods of Soil Analysis II, Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, WI, pp 595-624.
- Darrah PR. 1993. The rhizosphere and plant nutrition: a quantitative approach. *Plant Soil*. 155/156: 1-20.
- Djaenuddin D, Basuni, S Hardjowigeno, H Subagjo, M Sukardi, Ismangun, Marsudi Ds, N Suharta, L Hakim, Widagdo, J Dai, V Suwandi, S Bachri and ER Jordens. 1994. Land Suitability for Agricultural and Silviculture Plants. Laporan Teknis No. 7. Versi 1.0. April 1994. Center for Soil and Agroclimate Research, Bogor (in Indonesian).
- Dougan WK and Wilson AL. 1974. The absorptiometric determination of aluminum in water. A comparison of some chromogenic reagent and the development of an improved method. *Analyst* 99: 413-430.
- Eo J and T Nakamoto. 2006. Evaluation of root effect on soil organisms under different fertilization regimes by comparing rhizosphere and interrow soil in a wheat field. *Plant Root* 1: 3-9.
- Gahoonia TS, S Raza and NE Nielsen. 1994. Phosphorus depletion in the rhizosphere as influenced by soil moisture. *Plant Soil* 159: 213-218.
- Gee GW and JW Boulder. 1986. Particle size analysis. In: A Klute (ed). *Methods of Soil Analysis I, Physical and Mineralogy Methods*, 2nd edition. ASA, Madison, WI, pp. 383-412.
- Grinsted MJ, MJ Hedley, RE White and PH Nye. 1982. I. pH change and the increase in P concentration in the soil solution. *New Phytol* 91: 19-29
- Hasegawa T, E Purnomo, Y Hashidoko, M Osaki and G Rusmayadi. 2004. Grain yield and its variation of local rice varieties grown on acid sulphate soil in South Kalimantan. *Japanese J Crop Sci* 73: 220-221.
- Hinsinger P, GR Gobran, PJ Gregory and WW Wenzel. 2005. Rhizosphere geometry and heterogeneity arising from rootmediated physical and chemical processes. *New Phytol* 168 : 293-303.
- Houba VJG, I Novozamsky and E Temminghoff. 1994. Soil Analysis Procedures Extraction with 0.01 M CaCl_2 (Soil and Plant Analysis, Part 5A). Department of Soil Science and Plant Nutrition. Wageningen Agricultural University. The Netherlands.
- John MK. 1970. Colorimetric determination of phosphorus in soil and plant materials with ascorbic acid. *Soil Sci* 100: 214-220.

- Kempers AJ and A Zweers. 1986. Ammonium determination in soil extract by the salicylate method. *Comm in Soil Sci Plant Anal* 17: 715-723.
- Knudsen D, GA Peterson and PF Prafit. 1982. Lithium, sodium and potassium. In: AL Page, RH Miller and DR Keeney (eds), *Methods of Soil Analysis II, Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, WI, pp. 225-246.
- Li H, X Yang, GJD Kirk and A Dobbermann 2002. Root-induced of potassium in the rhizosphere of lowland rice (*Oriza sativa* L.) 17th WCSS, 14-21 August 2002, Thailand.
- Li X-L, E George and H Marschner. 1991. Phosphorus depletion and pH decrease at the root-soil and hyphae-soil interfaces of VA Mycorrhizal White Clover fertilized with ammonium. *New Phytol* 119: 397-404.
- Luster J, A Gottlein, B Nowack and G Sarret. 2009. Sampling, defining, characterising, and modelling the rhizosphere-the soil science tool box. *Plant Soil* 321: 457-482.
- McLean EO. 1982. Soil pH and lime requirement. In: AL Page, RH Miller and DR Keeney (eds). *Methods of Soil Analysis. II. Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, Wisconsin, pp 199-224.
- Olsen SR and LE Sommers. 1982. Phosphorus. In: AL Page, RH Miller and DR Keeney (eds), *Methods of Soil Analysis II, Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, Wisconsin, pp: 403-430.
- Purnomo E. 1996. Mineralisation and nitrification of soil under cereal crop. *PhD Thesis*. Charles Sturt University. Wagga, NSW.
- Purnomo E, M Sarwani, A Jumberi, A Mursyid, T Hasegawa, Y Hashidoko, T Shinano, S Honma and M Osaki. 2005. Phosphorus Nutrition of High Yielding Local Rice Varieties Grown without Fertilizer on Acid Sulphate Soil. *Soil Sci Plant Nutr* 51: 679- 681.
- Purnomo E, Y Hashidoko, T Hasegawa and M Osaki. 2010. Extreme high yield of tropical rice grown without fertilizer on acid sulfate soil in South Kalimantan, Indonesia. *J Trop Soils* 15 (1): 33-38.
- Rhoades JD. 1982a. Cation exchange capacity. In: AL Page, RH Miller and DR Keeney (eds), *Methods of Soil Analysis II, Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, WI, pp. 149-158.
- Rhoades JD. 1982b. Soluble salts. In: AL Page, RH Miller and DR Keeney (eds), *Methods of Soil Analysis II, Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, WI, pp. 167-180.
- Saleque MA. and GJD. Kirk 1995. Root-induced solubilisation of phosphate in the rhizosphere of lowland rice. *New Phytol* 129: 325-336.
- Thomas GW. 1982. Exchangeable cations. In: AL Page, RH Miller and DR Keeney (eds), *Methods of Soil Analysis II, Chemical and Microbiological Properties*, 2nd edition. ASA, Madison, WI, pp. 159-166.
- Yanai RD, H Majdi and BB Park. 2003. Measured and modelled differences in nutrient concentrations between rhizosphere and bulk soil in a Norway spruce stand. *Plant Soil* 257: 133-142.
- Yang JE, EO Skogley, BE Schaff and JJ Kim. 1998. A simple spectrophotometric determination of nitrate in water, resin and soil extract. *Soil Sci Soc Am J* 62: 1108-1115.
- Yeomans JC and JM Bremner. 1988. A rapid and precise method for routine determination of organic carbon in soil. *Comm in Soil Sci Plant Anal* 19: 1467-1476.