

The Influence of Organic and Inorganic Amendments on Phosphorus Chemistry in Two Acidic Soils of Southwestern Ghana

Owusu-Gyimah Victor¹, Klogo Phaniel Yao² and Gbefo Francis^{3*}

¹Department of Agriculture Engineering, Ho Technical University, P. O. Box HP 217, Ho, Ghana

²Department of Agricultural Science and Technology, Ho Technical University, P.O Box HP 217, Ho, Ghana

³Department of Science, Peki College of Education, P.O Box 14, Peki, Ghana.

*e-mail: francis.gbfo@pekicoe.edu.gh

Received 06 September 2024, Revised 28 October 2024; Accepted 02 January 2025

ABSTRACT

A study in southwestern Ghana compared the effects of organic and inorganic additives on P availability and related factors in two acidic soils, Ankasa and Abenia. Different amounts of P as KH_2PO_4 were applied: 0.067 g kg^{-1} for Abenia and 0.041 g kg^{-1} for Ankasa. Soil samples were treated with cow dung, *Chromolaena odorata*, and poultry droppings for six weeks to increase standard P requirement and neutralize exchangeable Aluminum. Data analysis was performed using GenStat (version 14). An analysis of variance (ANOVA) was conducted for the soil amendments, followed by Tukey's comparison test at a 5% significance level to identify significant differences among the soil amendments. The result showed that higher rates of organic amendments significantly increased pH, available P (Bray 1 and $\text{NaHCO}_3\text{-P}$), NaOH-P , and reduced exchangeable Al concentration. Poultry droppings and cow dung impact notably improved soil quality. At the same time, CaCO_3 had similar effects on soil pH. However, it did not significantly affect P availability or $\text{NaOH-extractable P}$. CaSO_4 and CaCO_3 had minimal impact on phosphorus distribution, suggesting that altering pH or exchangeable Al does not necessarily change P fractions. Poultry droppings, rich in P, could be a potential alternative to lime in enhancing P availability and reducing soil acidity.

Keywords: Acidic soil, exchangeable Aluminum, organic and inorganic amendments, phosphorus fractions

INTRODUCTION

The soil in southwestern Ghana, accounting for 10-15% of the country's land, is predominantly acidic despite favorable topography and hydrology, leading to low crop production (Ofori-Sarpong & Amankwah, 2019; Agegnehu *et al.*, 2021). Numerous investigations conducted over the years have consistently shown that adding phosphate fertilizer to tropical soils results in a substantial portion of the fertilizer being adsorbed onto colloidal surfaces in configurations that are not easily accessible to crops. This phenomenon leads to only a modest proportion, approximately 10-20%, of the applied fertilizer being effectively utilized by crops (Mabagala, 2022; Hanyabui, 2020).

The occurrence of sorbed phosphorus is common in weathered soils like Ultisols and Oxisols and volcanic soils with high amorphous compounds, primarily due to Al and Fe oxides and hydroxides

(Reed *et al.*, 2011; Amadou *et al.*, 2022). Contemporary approaches to soil management for mitigating P sorption include applying liming agents (such as CaCO_3), integrating P, or introducing organic amendments into soils with high P sorption capacity (Fan *et al.*, 2022). The latter approach is notably appealing in developing and developed nations, serving as a cost-efficient and environmentally friendly alternative to traditional inorganic P fertilizers and liming amendment (Huck *et al.*, 2014). Studies show conflicting results on the impact of organic amendments on P sorption and desorption in high P sorbing soils (Huck *et al.*, 2014; Serveshe *et al.*, 2015; Nobile *et al.*, 2020).

The results are due to the competition of organic acids, like citrate or malate, for sorption sites on Al and Fe, leading to the production of organic acids. Conversely, Yan *et al.* (2013) documented increased P sorption when incorporating organic amendments with low P content. Yusran (2018) also documented a positive relationship between soil organic matter content and P adsorption. The phenomenon is due to organic matter decomposition, where

microorganisms absorb P as lipids and nucleoproteins, making it accessible upon their death and decay. Research on phosphorus sorption in organic amendment-added soils is limited, lacking comparisons with inorganic amendments like CaCO_3 and CaSO_4 , which increase pH and enhance P availability (Servesh *et al.*, 2015; Melese & Yli-Halla, 2016).

However, the presence of CaSO_4 enhances the accessibility of P by displacing exchangeable Al and Fe while causing minimal alteration to the pH levels (Lizarralde *et al.*, 2021). This research aimed to compare the effects of organic and inorganic amendments on P desorption, pH, and exchangeable Al in two acidic soils of southwestern Ghana.

MATERIAL AND METHODS

Soil Samples and Amendment Materials

Two acid soils of agricultural importance, such as the Abenia and Ankasa of Southwestern Ghana, were chosen for the study. The soils are designated as Typic Hapludox by soil taxonomy (Soil Survey Staff, 1994), originate from biotite granite schist, and are found on the upper slope of the landscape. Surface soil (0-15 cm) was collected under a virgin tropical evergreen rainforest with an average rainfall of 1800 and 2000 mm per annum. The sampled soils were air-dried and sieved using a 2 mm sieve before analysis. Three organic amendments were used: cow dung, *Chromolaena odorata*, and poultry droppings. After air-drying, the materials underwent oven-drying at 60 °C for 24 hours before being ground and sieved through a 0.5 mm mesh. The sieved materials were then stored in white polyethylene bags before chemical analysis.

Laboratory Analysis

Soil pH was assessed in distilled water at a soil-to-solution ratio of 1:2.5 using a Sontex pH meter (Table 1). The total C content was quantified through

the wet oxidation technique developed by Walkley and Black (1934). The total N content in both soil and organic amendments was determined utilizing the Kjeldahl method. Additionally, extractable P levels were determined through the Bray 1 method (0.03M NH_3F and 0.25 M HCl solution), NaHCO_3 -P (Olsen *et al.*, 1954), and NaOH-P using a 0.1M NaOH solution. Total P was quantified by digesting the soil and organic samples with concentrated HNO_3 - HClO_4 until the solution became colorless. Subsequently, the ascorbic acid molybdate method developed by Watanabe and Olsen (1965) was employed following cooling and filtration. The color intensity was then assessed using a spectrophotometer set at a wavelength of 712 nm.

Exchangeable Al and acidity were extracted with 1M KCl solution (Barnhisel & Bertsch, 1982). Exchangeable bases were identified using 1 M NH_4OAc solution at a pH 7. The particle size analysis involved the complete breakdown of organic matter with a 1:1 mixture of soil and H_2O_2 . Subsequently, the soil particles were dispersed using a 0.01 M calgon (NaPO_3)₆ solution, followed by a five-minute mixing period with a motor mixer. Silt content was determined after five minutes using a hydrometer, while the clay fraction was assessed over 5 hours.

Incubation Studies

Based on the standard P requirement of these soils, P as KH_2PO_4 was initially incubated at rates of 0.067 g kg^{-1} and 0.041 g kg^{-1} in Abenia and Ankasa, respectively. The fertilizer was mixed thoroughly and then incubated for six weeks at room temperature (26 °C). Throughout the incubation period, samples were subjected to drying and wetting cycles after seven days, maintaining moisture content gravimetrically at 60% water holding capacity. Stirring was done every other day. Following the completion of the incubation period, the soil samples were air-dried and grounded for further laboratory testing.

Table 1. Chemical analysis of organic amendments.

| Organic Materials | Chemical analysis | | | | | | |
|---------------------|---|-----------------------|------|------|------|----------------------|------------------|
| | pH soil: H_2O 1:2.5 | cmol kg^{-1} | | | N % | Bray 1 (ppm) P | Organic C (%) |
| | | Ca | K | Mg | | | |
| Cowdung | 7.2 | 13.5 | 0.8 | 2.6 | 0.73 | 78 | 18.6 |
| Chromolaena Odorata | 7.5 | 28.4 | 1.82 | 5.86 | 2.23 | 80 | 20.23 |
| Poultry droppings | 8.2 | 30.36 | 1.65 | 5.43 | 2.35 | 264.8 | 23.73 |

Organic amendments (Cowdung, *Chromolaena odorata*, and Layer poultry droppings) were each added to the treated soil samples at an increasing rate of 0 (Control), 1% (Low), 2.5% (Medium), and 5% (High) (wt/wt). Another group of samples was treated with inorganic amendments (CaCO_3 and CaSO_4) at levels of 1.0 x (low), 1.5 x (medium), and 3.0 x (high) the quantity of CaCO_3 and CaSO_4 required to offset the exchangeable Al present in the soils. The amended soil samples were mixed thoroughly and made up to a moisture level of 60% water retention capacity using deionized water. All the samples were further incubated in triplicate at room temperature (26 °C) for six weeks.

The specimens were aerated on alternate days, and the water content was consistently monitored gravimetrically. After six weeks, the incubated samples underwent a process of air drying and grinding and were subsequently stored in zipped polyethylene bags for the determination of pH, exchangeable Al, available P (Bray 1, NaHCO_3 -P), and NaOH-P. All analyses were carried out in triplicates.

Statistical Analysis

Data were analyzed using analysis of variance (ANOVA) using Genstat (version 14). Post-hoc tests were employed to separate the means at a 5% probability level, while Microsoft Excel was used for the graphical representation of the data.

RESULTS

Impact of Organic and Inorganic Amendments on Soil Acidity Levels

The application of various organic amendments resulted in significant ($p < 0.05$) increases in soil pH compared to the control group. Cow dung exhibited the least impact on soil pH among the amendments in both types. The highest increase in pH was caused by *Chromolaena odorata* and poultry droppings. It was observed that increasing amounts of organic amendments caused a corresponding increase in pH in both soils. Interestingly, CaCO_3 and poultry droppings affected a similar magnitude of decreasing acidity in the two soils. In addition, increasing rates of CaSO_4 led to a proportional drop in pH. However, there was no statistically significant difference in the alteration of both soils. The order of effectiveness in decreasing soil acidity for organic and inorganic amendments was poultry droppings > CaCO_3 > *Chromolaena odorata* > cow dung > control > CaSO_4 (Figure 1).

Variation in Exchangeable Al^{3+} Concentration Due to Amendments

The impact of organic and inorganic amendments on exchangeable Al (Al^{3+}) in the two soils is shown in Table 2. The findings indicated that CaSO_4 had a minimal impact on decreasing the




Figure 1. Impact of organic and inorganic amendments on soil pH of two acidic soils of southwestern Ghana. ■ : 0%, ■ : 1%, ■ : 2.50%, ■ : 5%.

Table 2. Impact of organic and inorganic amendments on exchangeable Aluminum (Al^{3+}) in two acidic soils of Southwestern Ghana.

| Treatments / Levels | | | | | | | | | | | | | | | | |
|---------------------|-------------------------------------|---|-------------------|-------------------|---|-----|----|--|-------------------|-------------------|---|-------------------|----|---|------|----|
| Soils | Control (cmol+kg ⁻¹) | CaSO ₄ (cmol + kg ⁻¹) | | | CaCO ₃ (cmol + kg ⁻¹) | | | Cow dung (cmol + kg ⁻¹) | | | Chromolaena odorata (cmol + kg ⁻¹) | | | Poultry droppings (cmol + kg ⁻¹) | | |
| | 0% | 1x | 1.5x | 3x | 1x | 1.5 | 3x | 1% | 2.5% | 5% | 1% | 2.5% | 5% | 1% | 2.5% | 5% |
| Ankasa | 2.00 ^a | 2.00 ^a | 2.20 ^a | 2.50 ^b | nd | nd | nd | 1.20 ^c | 0.68 ^c | 0.30 ^c | 0.70 ^c | 0.50 ^d | nd | nd | nd | nd |
| Abenia | 0.81 ^a | 0.87 ^a | 0.88 ^a | 1.20 ^b | nd | nd | nd | 0.33 ^c | 0.25 ^c | nd | nd | nd | nd | nd | nd | nd |

LSD (P=0.05): P=0.093 in Abenia soil, P=0.16 in Ankasa soil, and = no exchangeable Al detected after six weeks of incubation. It means that letters bearing the same letters are not significantly different. X = time the quantity of CaCO₃ or CaSO₄ required to neutralize exchangeable Aluminum.

concentration of Al^{3+} in both soils, whereas CaCO₃ had the most significant effect, reducing Al^{3+} levels by up to 100%. The lowest level of CaCO₃ was optimal for ultimately reducing Al^{3+} in both soils. Generally, the addition of CaSO₄ did not result in a statistically significant ($p>0.05$) decrease in the concentration of Al^{3+} in the control group.

However, at the highest level (3x), CaSO₄ in both soils was found to increase the exchangeable Al significantly ($P<0.05$). The organic amendments significantly ($P<0.05$) reduce Al^{3+} compared to the control. Poultry droppings affected a 100% reduction in the concentration of Al^{3+} , followed by *Chromolaena odorata* (80-100%) and cow dung (70-100%). Increasing rates for all the organic amendments reduced the concentration of exchangeable Al (Table 2). The order of effectiveness in reducing the exchangeable Al followed the trend of poultry droppings > *Chromolaena odorata* > cow dung and control.

Impact of Organic and Inorganic Amendments on P Availability

In this study, Bray 1 and NaHCO₃-P were used as indices of P availability. The impact of CaCO₃ and CaSO₄ on availability P did not show a significant difference ($P>0.05$) from the control at lower application rates. However, at the highest level of CaSO₄ in both soils, there was a decrease in NaHCO₃-P. Whereas the observed decrease was significant ($P<0.05$) in Ankasa, it was insignificant in the Abenia soil. Abenia soil CaCO₃, at the highest level, caused a decrease in available P. However, there was not a statistically significant ($P>0.05$) difference in the effect.

On the other hand, the medium (1.5x) and highest (3x) levels of CaCO₃ in Ankasa soil exhibited a significant ($P<0.05$) increase from the control. Incorporating various organic amendments leads to a notable increase in the P availability in both soil




Figure 2. Impact of organic and inorganic amendments on P availability on two acidic soils of southwestern Ghana. ■ : 0%, ■ : 1%, ■ : 2.5%, ■ : 5%.

types. Unlike poultry droppings, the cow dung and *Chromolaena odorata* were ineffective in Abenia soil at the lower level of incorporation; however, in Ankasa soil, *Chromolaena odorata* at 1% was effective in increasing available P. Increasing rates of organic amendments increased available P. The order of effectiveness in increasing available P among the organic amendments followed the trend of poultry droppings > *Chromolaena odorata* > cow dung and control (Figure 2).

Impact of Organic and Inorganic Amendments on NaOH Extractable P

Soils amended with poultry droppings gave the highest increase in NaOH-P fraction, followed by *Chromolaena odorata* and cow dung. Increasing rates of the organic amendments led to a corresponding significant ($P < 0.05$) increase in NaOH-P fraction. In Ankasa soil, the lowest levels of *Chromolaena odorata* and cow dung resulted in a statistically significant ($P < 0.05$) elevation in NaOH-P levels. Applying inorganic amendments to the soils showed minimal impact on NaOH-P levels. Both CaSO_4 and CaCO_3 at low and medium levels did not significantly ($P > 0.05$) affect NaOH-P. However, it was observed that whereas CaCO_3 increased NaOH-P at the same level, CaSO_4 decreased NaOH-P. At the highest level, CaSO_4

significantly decreased NaOH-P in the Abenia soil but not in Ankasa (Figure 3). On the other hand, CaCO_3 significantly decreased and increased NaOH-P in Abenia and Ankasa, respectively.

DISCUSSION

The substantial increase in soil pH due to organic amendments in both soil types is significant for the reaction of P in acidic soils, as shown by the statistical significance level of $P < 0.05$. The elevation of pH levels leads to the precipitation of exchangeable Al and Fe, consequently diminishing the capacity for sorption of HPO_4^{2-} .

The rise in levels resulting from the use of organic additives can be credited to self-neutralization caused by the breakdown of C, the liberation of essential positively charged ions, or the release of hydroxide ions because of the decrease in Mn, Fe, and Al at oxygen-deprived microenvironments (Fageria & Nascente, 2014). High pH values of organic materials and high basic cations contribute to the rise in soil pH. The results are therefore consistent with the observation made by Huck *et al.* (2014). Alternatively, the rise in soil pH may be linked to generating OH^- ions through the ligand exchange process involving organic acids and hydroxyl ions of Al and Fe in the soil (Sokolova,




Figure 3. Impact of organic and inorganic amendments on NaOH extractable P on two acidic soils of southwestern Ghana. ■ : 0%, ■ : 1%, ■ : 2.50%, ■ : 5%.

2020). Nonetheless, a rise in soil pH by CaCO_3 could be due to the production of OH^- when it interacts with water in the soil. The resulting OH^- causes an increase in the pH by neutralizing the effect of the potential acidity in the soil (Mabagala, 2022). The decline in pH with increasing CaSO_4 may also be attributed to the formation of H_2SO_4 as CaSO_4 interacts with H_2O .

Under normal circumstances, the Ca ion is expected to produce more OH^- ions to neutralize the effect of H_2SO_4 , but the reverse is the case, indicating that more H_2SO_4 was produced such that the OH^- produced by CaSO_4 , even at the highest level of application was insufficient to neutralize it. The reduction of exchangeable Al because of organic amendments and CaCO_3 in the two soils may be partly a result of an increase in soil pH (Figures 1a and 1b). Huck *et al.* (2014) and Nobile *et al.* (2020) have also documented comparable findings. pH determines the activity of Al in each soil; at low pH, more Al^{3+} comes into the soil solution, while at high pH, little or no Al^{3+} comes into the solution (Mabagala, 2022). Alternatively, the decrease could also be ascribed to the formation of Al ions due to the OH^- ions being discharged from the substitution of ligands among organic anions and terminal hydroxyl of Fe and Al oxides or the binding of Al by organic compounds (Servesch *et al.*, 2015; Melese & Yli-Halla, 2016). Concerning CaSO_4 , the increase in Al^{3+} at the highest level may be attributed to the lowering of the pH in both soils (Figures 1 and 2). As pH decreases, more Al^{3+} comes into the soil solution, resulting in a higher Al^{3+} concentration. The impact of CaSO_4 on soil pH is consistent with findings reported by Zhao *et al.* (2022).

The available P indices (Bray 1 and NaHCO_3 -P) in both soil samples showed a notable rise in P levels at a significant level ($p < 0.05$) when the concentration reached 2.5% of organic amendments incorporation. The increase in available P by organic amendments may be credited to creating enduring complexes between Al and anions generated from the breakdown of organic substances. During the mineralization of organic amendments, organic acids like citrate, malate, or tartaric acid are liberated into soil solution and their anions complex Al, resulting in more P in solution (Philips, 2002; Sindhu *et al.*, 2022). The increase may also be a result of H_2PO_4^- anion by humate ions and the formation of a protective cover by humus around particles, leading to a decrease in the soil's phosphate sorption capacity (Huck *et al.*, 2014; Nobile *et al.*, 2020). Moreover, the increased level of accessible P due to organic amendments implies a more significant mineralization of P and the subsequent release of P

to interact with P on the adsorption sites of Al, ultimately resulting in an augmentation in P in the soil solution (Sarvest *et al.*, 2015; Sokolova, 2020). The increase may also be a result of the change in soil chemistry. These factors are recognized to significantly influence the availability of P and their absence or reduction in the soil system, especially Al^{3+} , and an increase in pH will increase available P. The high increase in P by poultry droppings in Abenia is at a 5% level, despite the pH values rising to about 7.51, possibly because poultry droppings are known to produce many citrates that can chelate exchangeable Al (Bauer, 2019).

Generally, the phosphorus-rich remains of poultry waste led to a notable rise in biologically accessible P1 (Bray 1 and NaHCO_3 -P). The findings are consistent with the results of Huck *et al.* (2014) and Nobile *et al.* (2020). CaSO_4 and CaCO_3 did not significantly impact the availability of phosphorus, despite CaCO_3 causing an increase at all levels except the highest in Albania. There was a slight decrease in this location, but it was not significant ($P > 0.05$). Conversely, there has been a notable reduction in the amount of P by CaSO_4 at the highest level in Ankasa soil. It may be attributed to the lowering of soil pH (Figures 1), thereby increasing Al^{3+} activity, as shown in Table 3.

Aluminum can sorb P, rendering P unavailable (Ofori-Sarpong & Amankwah, 2019; Agegnehu *et al.*, 2021). The decrease in available P in Abenia soil at the highest level of CaCO_3 may be due to precipitation of P by excess Ca in the soil solution enhanced by a rise in soil pH to 7.51, thereby reducing P availability. The NaOH extractant eliminates phosphorus and is not as strongly associated with P uptake by plants as NaHCO_3 -P (Obikoya, 2016). The fraction under consideration is linked to amorphous and crystalline Al and Fe-P through chemisorption (Linguist & Ruark, 2011). The rise in chemisorbed P fractions detected in both soil types, as indicated by NaOH-P, is probably due to the introduction of soluble inorganic P into the organic residue and the mineralization of P from the organic P that was added (Verma *et al.* 2005 & Sarvest, 2015).

CONCLUSIONS

Poultry manure affects a particular P fraction that holds significance in P sorption reactions. Incorporating organic residues with high total P content (like poultry droppings) modifies the distribution of P fractions (NaHCO_3 -p, Bray 1, and NaOH-P). Conversely, CaCO_3 and CaSO_4 showed limited to no influence on P fractions. Furthermore,

the rise in organic amendments applied to high P-fixing soil led to an increase in pH levels and a reduction in exchangeable Al content. This finding holds significant implications for the behavior of P in soil systems. The optimal level of organic amendments was 2.5%, suggesting that organic additives can be an alternative to traditional inorganic additives such as CaCO_3 to improve P availability in acidic soils.

ACKNOWLEDGMENTS

We are grateful to the Department of Soil Science, University of Ghana, Legon.

REFERENCES

- Agegehu, G., Amede, T., Erkossa, T., Yirga, C., Henry, C., Tyler, R., & Sileshi, G. W. (2021). A review of the extraction and management of acid soils for sustainable crop production system in the tropical agroecosystems. *Acta Agriculturae Scandinavica, Section B. Soil & Plant Science*, 71(9), 852–869.
- Amadou, I., Faucon, M. P., & Houben, D. (2022). New insights into sorption and desorption of organic phosphorus on goethite, gibbsite, kaolinite, and montmorillonite. *Applied Geochemistry*, 143, 105378.
- Bauer, P. J., Szogi, A. A., & Shumaker, P. D. (2019). Fertilizer efficacy of poultry litter ash blended with lime or gypsum as fillers. *Environments*, 6(5), 50.
- Fageria, N. K., & Nascente, A. S. (2014). Management of soil acidity of South American soils for sustainable crop production. *Advances in Agronomy*, 128, 221–275.
- Fan, B., Ding, J., Fenton, O., Daly, K., Chen, S., Zhang, S., & Chen, Q. (2022). Investigation of differential levels of phosphorus fixation in dolomite and calcium carbonate amended red soil. *Journal of the Science of Food and Agriculture*, 102(2), 740–749.
- Hanyabui, E., Obenga Pori, S., Agyei Frimpong, K., Atiah, K., Abindaw, T., Ali, M., & Byalebeka, J. (2020). Phosphorus sorption in tropical soils. *AIMS Agric. Food*, 5, 599–616.
- Huck, Y. C., Osumanu, H. A., & Nik, M. A. M. (2014). Improving phosphorus availability in an acid soil using organic amendments produced from agroindustrial wastes. *The Scientific World Journal*, 2014(1), 506356.
- Jackson, K. L. (1958). *Soil chemical analysis*. Prentice Hall.
- Linquist, B., & Ruark, M. (2011). Reevaluating diagnostic phosphorus tests for rice systems based on soil phosphorus fractions and field-level budgets. *Agronomy Journal*, 103(2), 501–508.
- Lizarralde, C. A., McDowell, R. W., Condron, L. M., Brown, J., & Whelan, M. (2021). Amending soils of different pH to decrease phosphorus losses. *Soil Research*, 60(2), 114–123.
- Mabagala, F. S. (2022). On the tropical soils: The influence of organic matter (OM) on phosphate bioavailability. *Saudi Journal of Biological Sciences*, 29(5), 3635–3641.
- Melese, A., & Yli-Halla, M. (2016). Effects of lime, wood ash, manure, and mineral P fertilizer applications on the inorganic P fractions and other selected soil chemical properties on acid soil of Farta District, Northwestern Highland of Ethiopia. *African Journal of Agricultural Research*, 11(2), 87–99.
- Nobile, C. M., Bravin, M. N., Becquer, T., & Pailat, J. M. (2020). Phosphorus sorption and availability in an andosol after a decade of organic or mineral fertilizer applications: Importance of pH and organic carbon modifications in the soil compared to phosphorus accumulation. *Chemosphere*, 239, 124709.
- Obikoya, O. (2016). *Changes in soil test phosphorus and phosphorus forms with continuous phosphorus fertilizer addition to contrasting prairie soils* (Master's thesis). Department of Soil Science, University of Manitoba, Canada. <http://hdl.handle.net/1993/31736>
- Ofori-Sarpong, G., & Amankwah, R. K. (2019). Acid drainage potential of rocks in Southwestern Ghana. *International Journal of Environmental Protection and Policy*, 7(1), 9.
- Olsen, S. R., Cole, C. U., & Watanabe, F. S. (1954). Dean, L. A. (1954). Estimation of available P in soil extraction with sodium bicarbonate. *Circular, U.S. Department of Agriculture*, 929.
- Phillips, I. R. (2002). Phosphorus sorption and nitrogen transformation in two soils treated with piggery wastewater. *Soil Research*, 40(2), 335–349.
- Reed, S. C., Townsend, A. R., Taylor, P. G., & Cleveland, C. C. (2011). Phosphorus cycling in tropical forests growing on highly weathered soils. In *Phosphorus in action: Biological processes in soil phosphorus cycling* (pp. 339–369).
- Sarvest, K., Ajaya, S., & Amid, G. (2015). Effect of organic amendments on availability of different chemical fractions of phosphorus. *Agricultural Science Digest*, 35(2), 83–88.
- Sindhu, S. S., Sehwat, A., & Glick, B. R. (2022). The involvement of organic acids in soil fertility, plant health, and environmental sustainability. *Archives of Microbiology*, 204(12), 720.
- Soil Survey Staff. (1994). *Keys to soil taxonomy* (5th ed.). United States Department of Agriculture, Soil Conservation Service.
- Sokolova, T. A. (2020). Low-molecular-weight organic acids in soils: Sources, composition, concentrations, and functions: A review. *Eurasian Soil Science*, 53, 580–594.

- Verma, S., Subehia, S. K., & Sharma, S. P. (2005). Phosphorus fractions in an acid soil continuously fertilized with mineral and organic fertilizers. *Biology and Fertility of Soils*, 41, 295–300.
- Walkley, A., & Black, I. A. (1934). An examination of the dictagraph method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37, 29–38.
- Watanabe, F. S., & Olsen, E. R. (1965). Test of an ascorbic method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil Science Society of America Proceedings*, 29, 677–678.
- Yan, X., Wang, D., Zhang, H., Zhang, G., & Wei, Z. (2013). Organic amendments affect phosphorus sorption characteristics in paddy soil. *Agriculture, Ecosystems & Environment*, 175, 47–53.
- Yusran, F. H. (2018). The relationship between phosphate adsorption and soil organic carbon from organic matter addition. *Journal of Tropical Soils*, 15(1), 1–10.
- Zhao, W. R., Shi, R. Y., Hong, Z. N., & Xu, R. K. (2022). Critical values of soil solution Al³⁺ activity and pH for canola and maize cultivation in two acidic soils. *Journal of the Science of Food and Agriculture*, 102(15), 6984–6991.