

# Phosphorus Extraction from Soil Constituents Using Bray P-1, Mehlich-1 and Olsen Solutions

Jamalam Lumbanraja, Sri Mulyani, Muhajir Utomo and Sarno

*Department of Soil Science Faculty of Agriculture University of Lampung, Jln. Prof. Soemantri Brodjonegoro, No. 1, Bandar Lampung 35145, Indonesia, e-mail: j.lumbanraja53@gmail.com, jamalam.lumbanraja@fp.unila.ac.id*

Received 02 January 2017/ accepted 01 May 2017

## ABSTRACT

The extraction methods using Bray P-1, Mehlich-1, and Olsen solutions are the most frequently used in equilibrium condition to estimate the available P in the soil constituents. However, each method can give some different values that may not describe the availability of soil P. Therefore, it is necessary to conduct a laboratory experiment to compare the three solutions in equilibrium and kinetics conditions for P release from soil colloids as a basic data for the future related to plant productions. The objective of this study was to compare the amounts of P release in equilibrium and kinetics conditions using Bray P-1, Mehlich-1, and Olsen solutions and the rate constants ( $k$ ) of P release from soil colloids using the three solutions of five soil constituents or treatments: (1) Soil (100% soil), (2) P-rock (100% phosphate rock), (3) compost (100% chicken manure compost), (4) soil + P-rock (75% soil + 25% phosphate rock), and (5) soil + P-rock + compost (50% soil + 25% phosphate rock + 25% chicken manure compost). Each extraction for each treatment was conducted in triplicate. The results indicate that the amounts of extracted P employing equilibrium condition in all treatments are significantly lower compared to those employing kinetics condition. The results also show that Mehlich-1 solution is the most powerful solution to extract P from soil constituents followed by Bray P-1 solution and Olsen solution, except that in the soil + phosphate rock + compost treatment in which the amount of P extracted using Mehlich-1 solution is the highest then followed by that of P extracted by Olsen solution and Bray P-1 solution both in equilibrium and kinetic conditions. The highest rate constants ( $k$ ) of the reaction in all treatments were found in the extractions using Mehlich-1 solution for desorbed P followed by that of using Bray P-1 solution and Olsen solution.

---

**Keywords:** Compost, equilibrium, kinetics, P extraction solutions, phosphate rock, soil

## ABSTRAK

Metode ekstraksi Bray P-1, Mehlich-1, dan Olsen adalah yang paling sering digunakan dalam kondisi kesetimbangan untuk memperkirakan fosfor (P) yang ada dalam konstituen tanah. Namun masing-masing metode tersebut dapat memberikan nilai berbeda yang mungkin tidak menggambarkan ketersediaan P dalam tanah. Oleh karena itu, perlu dilakukan percobaan laboratorium untuk membandingkan ketiga larutan dalam kondisi kesetimbangan dengan kinetika untuk pelepasan P dari koloid tanah sebagai data dasar untuk masa datang terkait dengan produksi tanaman. Tujuan penelitian ini adalah untuk membandingkan jumlah P terekstrak dalam kondisi ekuilibrium dan kinetika dengan menggunakan larutan pengeksrak Bray P-1, Mehlich-1, dan Olsen dan konstanta kecepatan ( $k$ ) pelepasan P dari koloid tanah dengan menggunakan tiga larutan dengan perlakuan: (1) Tanah (100% tanah), (2) P-rock (100% batuan fosfat), (3) kompos (100% kompos kotoran ayam), (4) tanah + P-rock (50% tanah + 25% batuan fosfat + 25% kompos kotoran ayam) yang diekstraksi dalam tiga ulangan. Hasil penelitian menunjukkan bahwa jumlah P yang diekstraksi dalam kondisi ekuilibrium pada semua perlakuan secara nyata lebih rendah dibandingkan dengan menggunakan kinetika. Hasil penelitian juga menunjukkan bahwa larutan Mehlich-1 adalah yang paling banyak mengekstrak P dari konstituen tanah yang diikuti oleh larutan Bray P-1 dan paling sedikit adalah larutan Olsen, kecuali pada perlakuan tanah + batuan fosfat + kompos dimana jumlah P yang diekstraksi menggunakan larutan Mehlich-1 adalah yang tertinggi kemudian diikuti oleh P yang diekstraksi dengan larutan Olsen dan yang terendah adalah larutan Bray P-1 baik dalam kondisi kesetimbangan maupun kinetika. Konstanta kecepatan ( $k$ ) tertinggi dalam semua perlakuan ditemukan dengan menggunakan larutan Mehlich-1 untuk ekstraksi P diikuti larutan Bray P-1 dan yang paling rendah adalah dengan menggunakan larutan Olsen.

---

**Kata kunci:** kompos, ekuilibrium, kinetika, larutan ekstraksi P, batuan fosfat, tanah

## INTRODUCTION

Phosphorus (P) deficiency is one of the limiting factors to support the soil productivity in a humid tropical region in which the soil parent materials have undergone further weathering and leaching (Lumbanraja 1995; Lumbanraja *et al.* 1982; Oelkers and Valsami-Jones, 2008). The low pH of the soils can be an indication of high P retention by oxide of iron (Fe) and aluminum (Al), resulting in lower P availability to plants (Lumbanraja *et al.* 1981; Lumbanraja *et al.* 1994; Arai and Sparks 2007; De Araújo *et al.* 2015). The works have been done to improve the availability of P in soils, such as adding phosphate rock and manufacturing phosphate and organic fertilizers (Lumbanraja and Utomo 1996; Lumbanraja *et al.* 1995; Vetterlein *et al.* 1999; Lindsay *et al.* 1989; Gorgin *et al.* 2011). Natural phosphate rock is the primary raw material used in the manufacture of phosphate fertilizers. Natural phosphate rock comes from rocks containing mineral apatite that contains quite high amount of  $P_2O_5$  (Lindsay *et al.* 1989; Wahida *et al.* 2007). On the other hand, prediction of the amount of  $P_2O_5$  in soil, organic materials and phosphate rocks in general depends on the extraction methods employed using different kinds of extractant solutions (Gartley *et al.* 2002; Kweon *et al.* 2015).

Prediction of the availability of P in soil for any kinds of cultivated plants has to use appropriate methods of soil analysis. Phosphate availability is usually measured as a function of chemical equilibrium that is controlled by solubility and rate limited process (Sparks 1989; Lindsay *et al.* 1989; Arai and Sparks 2007; Kweon *et al.* 2015). Most methods used for the determination of available P attempt to quantify soluble P using different extractant solutions, but few of them related to P supply rates that are relevant to plant uptake (Ziadi *et al.* 2001; Lynch 2011; Meetei *et al.* 2015). Test methods for soil constituents including organic and phosphate rocks do not measure the amount of available P to a certain crop, but extract a portion of P from the soil constituents that is related to plant available P (Devau 2010; Lynch 2011; Rouached *et al.* 2010; Babana and Antoun 2006; Kweon *et al.* 2015). The amount of P extracted using different kinds of methods has to be related to variety of crop production, but the correlation will not always be transferable between crops and soils (Morel *et al.* 2014; Sing *et al.* 2005; Lumbanraja and Utomo 1996). It is because the immediately available P pool is constantly replenished through dissolution or desorption of less available P and through the mineralization of P in soil constituents such as

weathering of soil and phosphate rocks and decomposition of organic matter (Ziadi *et al.* 2001) and the available P for plant is strongly time dependent (van der Zee *et al.* 1988; van der Zee *et al.* 1986; Vetterlein *et al.* 1999).

There are several methods that are often used in the determination of soil P, namely Bray P-1, Mehlich-1, and Olsen. The most commonly used methods are the alkaline bicarbonate method (Olsen *et al.* 1954; Wolf and Baker 1985) and the acid ammonium fluoride extraction (Bray and Kurtz 1945; Wolf and Baker 1985) with various modifications. The bicarbonate method using  $NaHCO_3$  solution buffered at pH 8.5 (Olsen *et al.* 1954; Wolf and Baker 1985) has been used successfully in a wide range of acid to alkaline soils, especially in alkaline soils, while the acid ammonium fluoride extraction (Bray and Kurtz 1945) has been widely used on acid and neutral soils. Both methods are purely chemical soil tests that cannot be interpreted to plant function (Wolf and Baker 1985; Tiessen and Moir 2008). Various soil samples are commonly used for P extraction using Mehlich-1 method, expressing an excellent method to test the versatility of soil P extractant (Mehlich 1972). Mehlich-1 worked well in predicting plant-available P in acidic coastal plain soil (pH less than 6.5) with low cation exchange capacity ( $CEC < 10 \text{ cmol kg}^{-1}$ ) and low organic matter content ( $< 5\%$ , however, Mehlich-1 was not always adequate in predicting plant-available P in soils with high pH and CEC and significant accumulations of iron and aluminum such as in tropical Ultisols. Mehlich-1 method, which is suitable applied on acidic soils, can extract P in the form of Al-P, Fe-P and Ca-P (Meetei *et al.* 2015).

Phosphorus extraction methods usually take place in an equilibrium state to determine the available P for plant uptake. However, the equilibrium reaction is found less corresponding to availability of P for plant uptake (Meetei *et al.* 2015). The amount of P being desorbed from the colloid of soil constituents or phosphate rocks in the equilibrium reaction will be limited by the P concentration in the solution (pseudo solubility product -  $K_{sp}$ ). Meanwhile, the "true" reaction of P release into soil solution is an open reaction in which the reaction product of P loses through plant uptake or leaching in the soil systems (Koopmans *et al.* 2004; Maguire and Sims 2002; van der Zee *et al.* 1988; Torrent and Delgado 2001; Tipping *et al.* 2014). The desorbed P from soil colloids in the field soil systems will not be limited by the  $K_{sp}$  such as in the equilibrium condition but rather the released P will be absorbed by plant roots or reacted into a different form (Tiesen and Moir 2008; Sparks

1989; Schoumans 2013). Therefore, the kinetics approaches will be more reliable to express the availability of P in the soil reaction in describing the availability of P for plants (Lookman *et al.* 1995; Lookman *et al.* 1996; Lynch 2011; Nafiu 2009; van der Zee *et al.* 1986). The kinetics P can describe P release from soil colloids or phosphate rocks continuously using Bray P-1, Mehlich-1, and Olsen solutions. In the kinetics reaction, the cumulative product of P, which is extracted from the soil reactor, is expected to be higher than that in the equilibrium reaction. The purposes of the study were (1) to compare the amount of phosphate extracted by Bray P-1, Mehlich-1 and Olsen solutions in equilibrium and kinetics conditions, and (2) to compare the rate constants (kinetics) of P release using Bray P-1, Mehlich-1 and Olsen solutions in determining extractable P in a soil sample, compost and phosphate rock.

**MATERIALS AND METHODS**

The soil constituents used in this study were phosphate rock from Egypt, organic material (*i.e.* compost from chicken egg production) and a soil sample taken from the Experimental Station of University of Lampung in Natar, South Lampung District. The extractants used for determining the available P were Bray 1 (0.03 M NH<sub>4</sub>F + 0.025 M HCl), Mehlich-1 (0.05 M HCl + 0.0124 M H<sub>2</sub>SO<sub>4</sub>), and Olsen (0.5 M NaHCO<sub>3</sub> buffered at pH 8.5) solutions (Bray and Kurtz 1945; Mehlich 1972; Olsen *et al.* 1954). The P desorption reaction was conducted using a sequential extraction using batch techniques (Sparks 1989; Tiessen and Moir 2008) for the samples in the treatments of B<sub>1</sub> (100% soil), B<sub>2</sub> (100% phosphate rock), B<sub>3</sub> (100% compost), B<sub>4</sub> (75% soil + 25% phosphate rocks), B<sub>5</sub> (50% soil + 25% phosphate rock + 25% organic material). The amount of P extracted in the equilibrium condition of all treatments (B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>4</sub>, and B<sub>5</sub>) were determined using Bray P-1, Mehlich-1 and Olsen solutions (Bray and Kurtz 1945; Mehlich 1972; Olsen *et al.* 1954). On the other hand, the amount of P extracted in kinetics condition was determined using a sequential extraction (Tiessen and Moir 2008) followed by the cumulative extraction of day 1 + day 2 + day 4 + day 8 + day 16 + day 32 + day 64. About 1.5 g of samples was weighed in triplicate and then placed into a bottle, then 15 ml of P extractant solutions (Bray P-1, Mehlich-1, and Olsen) was added to the respective bottle. The suspensions were shaken using a shaker with the speed of 500 rpm for 30 minutes. The soil constituent suspensions were centrifuged at a speed of 5000

rpm for 10 minutes, then filtered using a filter paper to separate the solids and the clear solutions. The solids were subsequently extracted back using fresh extractant solutions of Bray P-1, Mehlich-1 and Olsen, respectively. The clear solution from each sequential extraction was used to determine the concentration of P by using a spectrophotometer at a wavelength of 720 nm (Murphy and Riley 1962; Thom and Utomo 1991).

The data were tabulated and interpreted based on the results in equilibrium and kinetics conditions. The following kinetics equations that describe the P release rate of the reactants (soil, phosphate rock and organic material) according to pseudo first order kinetics equation (Sparks 1989).

$$X - P_0 \rightarrow X - P_t \rightarrow P_t \dots \dots \dots (1)$$

in which  $X - P_0$  is the fraction of P at zero time,  $X - P_t$  is the fraction of P in the colloid at time  $t$ , and  $P_t$  is the accumulation of P fraction removed from the soil reactor at time  $t$ .

$$dP/dt = kP^n \dots \dots \dots (2)$$

in which  $P$  is the fraction of P in the colloid ( $(P_0 - P_{ex}) / P_0$ ) and  $n$  is the order of the reaction, and  $t$  is time (days),  $P_{ex}$  is the extracted P at time  $t$ , and  $k$  is the rate constant. For the pseudo first order kinetics:

$$dP/dt = kP \dots \dots \dots (3)$$

$$dP/P = kdt \dots \dots \dots (4)$$

by integration of equation (4) then,

$$\ln P_0 - \ln P_t = kt \dots \dots \dots (5)$$

rearranged

$$\ln P_t = \ln P_0 - kt \dots \dots \dots (6)$$

When  $\ln P_t$  is plotted as a function of  $t$  (day), it is a linear equation to create the kinetics figure of pseudofirst order to determine the rate constant  $k$ .

**RESULTS AND DISCUSSION**

The chemical properties of soil sample, P content in the phosphate rock, and P content in the organic material are presented in Table 1. The soil (Ultisol) from the Experimental Station of the University of Lampung in Natar has low fertility status as indicated by low pH (acid soil), low P-potential, very low available-P and exchangeable-

Table 1. Chemical properties of Ultisol from Natar, P content in the phosphate rock and P content in the organic matter used in the study.

Chemical properties	Value	Criteria*
pH H <sub>2</sub> O	5.05	Acid
pH KCl	4.82	Acid
Total-N (%)	0.06	Very low
Available-P(Bray P-1) (mg kg <sup>-1</sup> )	2.66	Very low
Exchangeable-K (cmol kg <sup>-1</sup> )	0.24	Low
Total-C (%)	0.43	Very low
CEC (cmol kg <sup>-1</sup> )	5.73	Low
P <sub>2</sub> O <sub>5</sub> of soil by 25% HCl (mg kg <sup>-1</sup> )	197.64	low
P <sub>2</sub> O <sub>5</sub> of phosphate rock by 25% HCl (%)	25.87	High
P <sub>2</sub> O <sub>5</sub> of organic by 25% HCl (%)	3.14	High

\*(Soil Research Institute 2005).

K, and very low total nitrogen and total-C. The low content of the nutrients in the soil is probably due to the soil has undergone further weathering and most of organic materials in the soil have been lost due to leaching processes and intensive decomposition. The potential P contents of the phosphate rock and organic material extracted using 25% HCl are considered as high, *i.e.* 25.87% (P<sub>2</sub>O<sub>5</sub>) and 3.14% (P<sub>2</sub>O<sub>5</sub>), respectively.

#### Comparison of the Amounts of Extracted P in Equilibrium and Kinetics Conditions

The cumulative curve of P concentrations extracted from soil sample using different extractant solutions of Bray P-1, Mehlich-1, and Olsen is presented in Figure 1, and the cumulative P concentrations determined using the three solutions in equilibrium and kinetics conditions (sequential

extraction) are presented in Table 2. The cumulative extracted P from soil increases as a function of extraction time (days) (Figure 1). It showed that the Mehlich-1 solution extracts the highest amount of P, followed by Bray P-1 and Olsen solutions. The amount of P extracted from the soil by Bray P-1 solution in equilibrium condition (*i.e.* 2.84 mg kg<sup>-1</sup>) is about nine times less than that of in kinetics condition (*i.e.* 25.98 mg kg<sup>-1</sup>). Similarly, the amount of P extracted from the same soil by Mehlich-1 solution in equilibrium condition (*i.e.* 4.64 mg kg<sup>-1</sup>) is about eight times less than that in kinetics condition (*i.e.* 33.93 mg kg<sup>-1</sup>) and the amount of P extracted by Olsen solution in equilibrium condition (*i.e.* 1.65 mg kg<sup>-1</sup>) is about 14 times less than that in kinetics condition (*i.e.* 21.70 mg kg<sup>-1</sup>).

Similar trends as those in the soil sample were found for all other treatments (100% phosphate rock,




Figure 1. Cumulative extracted P from soil as a function of time (t) using Bray P-1, Melich-1 and Olsen  
 ..... —◆— : Bray, ..... —■— : Mehlich, -▲- : Olsen.


Table 2. Comparison of the amounts of P in each treatment extracted using Bray P-1, Mehlich-1, and Olsen solutions in equilibrium and kinetics conditions.

Treatment	Extractant Solution	Extracted P (mg kg <sup>-1</sup> )		t-calculated	t-Table	
		Equilibrium	vs Kinetics		(0.05)	(0.01)
Soil	Bray P-1	2.84	25.98	7.85*		
	Mehlich-1	4.64	33.93	11.89*	6.31	12.71
	Olsen	1.55	21.70	11.58*		
P-rock	Bray P-1	18.94	236.19	14.55**		
	Mehlich-1	6,408.28	49,970.73	23.85**		
	Olsen	9.28	96.61	16.21**		
Compost	Bray P-1	2,979.21	6,467.87	7.98*		
	Mehlich-1	3,612.20	29,155.17	17.14**		
	Olsen	1,880.67	6,959.33	8.14*		
Soil+P-rock	Bray P-1	23.29	235.84	8.37*		
	Mehlich-1	4,822.87	61,176.83	54.42**		
	Olsen	3.18	47.28	17.48**		
Soil+P-rock+ Compost	Bray P-1	34.82	356.60	14.76**		
	Mehlich-1	5,622.88	49,978.46	33.82**		
	Olsen	63.91	487.65	13.81**		

\*= significantly different at 5% significance level; \*\*= significantly different at 1% significance level.

100% compost, 75% soil + 25% phosphate rock and 50% soil + 25% phosphate rock + 25% compost) (Table 2). The amounts of extracted P employing equilibrium condition in all treatments are significantly lower compared to that in kinetics condition. Especially, the amounts of extracted P employing equilibrium condition in the presence of phosphate rock are highly significantly (\*\* significant at 1%) lower compared to that in the kinetics condition, except that in the mixture of soil and phosphate rock of the extraction of Bray P-1 solution in which the difference on the amount of extracted P is significant (\* significant at 5%).

It also indicated that the amount of extractable P in each treatment extracted by Bray P-1, Mehlich-1, and Olsen solutions in equilibrium condition is approximately 10 times lower than that of extracted P in the kinetics condition. The reason for these results can be explained as the following reactions (7) and (9):




In which  $P_t$  is not removed from the soil reactor, then:

$$K_{seq} = (X-P_t) / (P_t) \dots \dots \dots (8)$$

In the equilibrium condition (7), the product of extracted  $P_t$  in the solution controls the equilibrium

constant of “extractability product” ( $K_{seq}$ ) and therefore when the  $P_t$  product in the solution meets the  $K_{seq}$  (8), then the P from colloids will not be removed any more or the reaction stops (equilibrium), while in kinetics, the following reaction takes place (9):



In which  $P_t$ (9) is removed from the soil reactor and the extracted solution is renewed and therefore, the  $P_t$  product in the solution can not control the  $K_{seq}$  (8), then the P from colloids ( $X-P_t$ ) can be removed or extracted continuously or the reaction can still continue until asymptotically close to the quantity of potential P (referring to Figure 1 and Table 1, especially the quantity of  $P_2O_5$  extracted by 25% HCl that is considered as potential extractable P) that can be extracted from the colloids.

**Comparison of the Amounts of Extracted P Employing Bray P-1, Mehlich-1 and Olsen Solutions in Equilibrium and Kinetics Conditions**

The results showed that Mehlich-1 solution is the most powerful solution to extract P from the soil constituents followed by Bray P-1 solution and Olsen solution, except that in the soil + phosphate rock + compost treatment in which the amount of P extracted using Mehlich-1 solution is the highest

Table 3. Comparison of extracted P employing Bray P-1, Mehlich-1, and Olsen solutions in equilibrium and kinetics conditions.

Extractant solution	t-calculated of extracted P		t-table	
	Equilibrium	Kinetics	0.05	0.01
Mehlich-1 vsBray P-1	74.85**	94.23**	6.31	12.71
Bray P-1 vs Olsen	51.70**	3.01 <sup>ns</sup>		
Mehlich-1 vs Olsen	71.73**	136.10**		

ns= not significantly different \*= significantly different at 5% significance level; \*\*= significantly different at 1% significance level

followed by that in the Olsen solution and Bray P-1 solution both in equilibrium kinetics conditions (Table 2). The results in Table 3 also showed that the amounts of P extracted using Mehlich-1 solution are highly significantly (\*\* significant at 1%) higher compared to that of using Bray P-1 and Olsen solutions both in equilibrium and kinetics conditions. In addition, the amounts of P extracted using Bray P-1 solution are highly significantly (\*\* significant at 1%) higher compared to that of using Olsen solutions in equilibrium condition, but the amounts of P extracted using Bray P-1 and Olsen are not significantly different in the kinetics condition. One of the reasons could be related to the acidity of the solutions. Mehlich-1 solution consists of two acid solutions (0.05 M HCl + 0.0125 M H<sub>2</sub>SO<sub>4</sub>) that are highly acidic (Mehlich 1972); Bray P-1 solution consists of 0.025 M HCl + 0.03 M NH<sub>4</sub>F in which the F<sup>-</sup> ion will be as powerful as H<sub>2</sub>PO<sub>4</sub><sup>-</sup> and HPO<sub>4</sub><sup>2-</sup> for anion exchange (Bray and Kurtz, 1945); while Olsen solution consists of 0.5 M NaHCO<sub>3</sub> buffered at pH 8.5, which is more appropriate for calcareous soils (Olsen *et al.* 1954). The non significant difference between the amounts of P extracted using Bray P-1 and Olsen solutions in kinetics condition is not clear. The trends of the amounts of P extracted

using the two solutions are not consistent. For example, the amounts of P extracted using Bray P-1 solution are higher compared to that of Olsen solution in the treatments of (1) soil, (2) phosphate rock and (3) soil + phosphate rock. On the other hand, the amounts of P extracted using Olsen solution are higher compared to that of Bray P-1 solution in the treatments of (1) compost and (2) soil + phosphate rock + compost. The interesting result is that the amounts of P extracted using Olsen solution are consistently higher compared to that of using Bray P-1 in the treatment of soil + phosphate rock + compost both in equilibrium and kinetics conditions.

#### Kinetics of P Release from Soil Constituents Employing Bray P-1, Mehlich-1, and Olsen Solutions

Figure 2 indicates the fraction of P remained in the colloids at time *t*, showing the lines of linear curves started from the steeper (higher) slopes at the beginning of the reactions followed by the lower asymptotic slopes at the later reactions. In kinetics reaction, P in soil solution as the product of P release is removed from the soil reactor, the rate of the P release from colloids will be determined by the




Figure 2. P-fraction in soil extracted using Bray P-1, Melich-1 and Olsen solutions as a function of time

—◆— : Bray, ...■... : Mehlich, -▲- : Olsen.

Table 4. Linear equations of P kinetics in each treatment determined using Bray P-1, Mehlich-1 and Olsen solutions.

Treatment	Extractant Solution	Linier equation	k	r <sup>2</sup>
Soil	Bray P-1	y = -0.0264x - 0.143	-0.0264	0.961
	Mehlich-1	y = -0.0472x - 0.138	-0.0472	0.986
	Olsen	y = -0.0196x - 0.087	-0.0196	0.959
P-rock	Bray P-1	y = -0.0174x - 0.001	-0.0174	0.913
	Mehlich -1	y = -0.0486x - 0.159	-0.0486	0.985
	Olsen	y = -0.0125x - 0.001	-0.0125	0.931
Compost	Bray P-1	y = -0.0052x - 0.095	-0.0052	0.695
	Mehlich -1	y = -0.0725x - 0.267	-0.0725	0.976
	Olsen	y = -0.0054x - 0.081	-0.0054	0.827
Soil + P-rock	Bray P-1	y = -0.0285x - 0.001	-0.0285	0.923
	Mehlich -1	y = -0.0487x - 0.158	-0.0487	0.986
	Olsen	y = -0.0215x - 0.001	-0.0215	0.911
Soil + P-rock+ Compost	Bray P-1	y = -0.0101x - 0.001	-0.0101	0.914
	Mehlich -1	y = -0.0477x - 0.150	-0.0477	0.988
	Olsen	y = -0.0082x - 0.002	-0.0082	0.901

Note: k = rate constant of P; r<sup>2</sup> = coefficient determination; y= ln P<sub>t</sub>; x= time t (day)

concentration of P in the colloids and the energy involved such as acidity and ionic strength of particular solution to dissolve or extract P from the soil colloids (Wolf and Baker 1985; Tiessen and Moir 2008; Sparks 1989).

The data in Table 4 are the pseudo first order of the reaction employing the linear equation of  $\ln P_t = \ln P_o - kt$  (6), in which the concentration of P<sub>t</sub> is expressed in the fraction of P in soil colloids as a function of time t (day) and the slope k represents the rate constant of the reaction (Nafiu 2009; Lookman *et al.* 1995; Sparks 1989). Figure 3 represents the  $\ln [P]_t$  as a function of time in the pseudo first-order reaction of P desorption from soil,

Figure 4 represents that from phosphate rock and Figure 5 represents that from compost.

The equations of the  $\ln [P]_t$  as a function of time and the rate constants that are represented by the slopes of the linear equations for all treatments are presented in Table 4. The highest rate constants (k) of the reactions in all treatments are found in the extractions using Mehlich-1 solution for desorbed P followed by that of using Bray P-1 solution and that of using Olsen solution. The reasons for this finding could be due to the Mehlich-1 solution is the most powerful solution to release P from colloids since Mehlich-1 solution is the strongest acidic solution compared to that of Bray P-1 and Olsen




Figure 3.  $\ln [P]_t$  as a function of time on the pseudo first-order reaction of P desorption from soil. —◆—: Bray, ...■...: Mehlich, -▲- : Olsen.




Figure 4.  $\ln [P]_t$  as a function of time on the pseudo first-order reaction of P desorption from phosphate rock.

—◆— : Bray, ...■... : Mehlich, -▲- : Olsen.




Figure 5.  $\ln [P]_t$  as a function of time on the pseudo first-order reaction of P desorption from compost.

—◆— : Bray, ...■... : Mehlich, -▲- : Olsen.

solutions, therefore, Mehlich-1 solution has more power to extract P from Al-P, Fe-P, and Ca-P from soil colloids, and the Mehlich-1 extraction is more suitable for acid soils (pH < 6.5) with CEC less than 10 cmol kg<sup>-1</sup> and organic matter content less than 5% (Wolf and Baker 1995). Mehlich-1 solution also has the highest ionic strength [I] compared to Bray P-1 and Olsen solutions (Evangelou 1998).

## CONCLUSIONS

The amounts of extracted P employing equilibrium condition in all treatments are significantly lower compared to that in kinetics condition. The results also showed that Mehlich-1 solution is the most powerful solution to extract P followed by Bray P-1 solution and Olsen solution, except that in the soil + phosphate rock + compost treatment in which the amount of P extracted using Mehlich-1 solution is the highest followed by Olsen solution and Bray P-

1 solution both in equilibrium and kinetics conditions. The highest rate constants ( $k$ ) of the reaction in all treatments were found in the extractions using Mehlich-1 solution for desorbed P followed by that of using Bray P-1 solution and Olsen solution. Further researches are expected to study the relationships of these findings to any kinds of crop productions in the field.

## REFERENCES

- Arai Y and DLSparks. 2007. Phosphate reaction dynamics in soils and soil minerals: a multiscale approach. *Adv Agron* 94: 135-179.
- Babana AH and H Antoun. 2006. Effect of Tilemsi phosphate rock-solubilizing microorganism on phosphorus uptake and yield of field-grown wheat (*Triticum aestivum* L.) in Mali *Plant Soil* 287: 51-58.
- Bray RH and LT Kurtz. 1945. Determination of total organic and available forms of phosphate in soils. *Soil Sci* 59: 39-45.



- De Araújo L, E Sampaio and M De Araújo. 2015. Phosphorus desorption from Fe and Al oxides mediated by soil microorganisms. *Commun Soil Sci Plant Anal* 46: 633-640.
- Devau N, E Le Cadre, P Hingsinger and FGérard. 2010. A mechanistic model for understanding root-induced chemical changes controlling phosphorus availability. *Ann Bot (Lond)* 105: 1183-1197.
- Evangelou VP. 1998. Environmental Soil and Water Chemistry: Principle and Application, John Wiley & Sons Inc. New York, 564 p.
- Gartley KL, JT Sims, CT Olsen and P Chu. 2002. Comparison of soil test extractants used in mid-Atlantic United States. *Commun Soil Sci Plant Anal* 33: 873-895.
- Gorgin N, M Fekri and L Sadegh. 2011. Impact of organic-matter application on phosphorus-desorption kinetics in two agricultural soils in Southeastern Iran. *Commun Soil Sci Plant Anal* 42: 514-527.
- Koopmans GF, WJ Chardon, P de Willigen and WH van Riemsdijk. 2004. Phosphorus desorption dynamics in soil and the link to a dynamic concept of bioavailability. *J Environ Qual* 33:1393-1402.
- Kweon G, E Lund, C Maxton, W Lee and D Mengel. 2015. Comparison of soil phosphorus measurements. *T Am Soc Agric Biol Eng* 58: 405-414.
- Lindsay WL, PLG Vlek and SH Chien. 1989. Phosphate minerals. In: JB Dixon and SB Weed (eds). Minerals in Soil Environment, 2<sup>nd</sup> Ed. Soil Science Society of America, Madison, WI, pp. 1089-1130.
- Lookman R, D Freese, R Merckx, K Vlassak and WH van Riemsdijk. 1995. Long-term kinetics of phosphate release from soil. *Environ Sci Technol* 29:1569-1575.
- Lookman R, K Jansen, R Merckx and K Vlassak. 1996. Relationship between soil properties and phosphate saturation parameters a transect study in northern Belgium. *Geoderma* 69: 265-274.
- Lumbanraja J. 1995. Sorption behavior of phosphorus in acid soil of Ultisols: Effects of phosphorus sources and urea. *J Pen Pengb Wil Lahan Kering* 15:1-17 (in Indonesian).
- Lumbanraja J, T Tadano, M Ninaki, K Oya, T Yoshida, M Utomo and AD Sitorus. 1981. Nutritional limitation factors for rice and corn growth on Podsolc Soils in Lampung Province. In: Y Asada (eds). Cooperation Research between DGHE and JSPS. Ehime University, Japan, pp.118-124.
- Lumbanraja J, T Tadano, M Ninaki, K Oya, T Yoshida, M Utomo and AD Sitorus. 1982. The effect of nitrogen, phosphorus, and potassium fertilizers on cassava growth on "Bandarjaya Podsolc Soil" and "Gedung Meneng Latosol Soil". In: Y Asada (eds) Cooperation Research between DGHE and JSPS. Ehime University, Japan, pp.112-122.
- Lumbanraja J and M Utomo. 1996. Phosphorus availability for rice and P-sorption behavior in soil affected by P-source and N-fertilizer in an acid soil of Sumatera. IVth International Symposium on Plant-Soil Interactions at Low pH. Brazil, March 17-24.
- Lumbanraja J, M Utomo and B Sitio. 1995. Sorption characteristics of soil phosphorus in the three different tillages and nitrogen fertilization in the experimental plot of a long-term experiment plots (in Indonesian). In: M Utomo, IH Utomo, FX Susilo, J Lumbanraja (eds) Proceedings of the Fifth National Seminar of Conservation Tillage. Bandar Lampung Indonesia pp. 69-77.
- Lumbanraja J, S Djuniwati and M Murad. 1994. The influence of particle size and the type of iron concretions and compost on the availability of phosphorus in the clay fraction of an Ultisol and an Alfisol (in Indonesian). *J Pen Pengb Wil Lahan Kering* 14:12-27.
- Lynch JP. 2011. Root phenes for enhanced soil exploration and phosphorus acquisition: tools for future crops. *Plant Physiol* 156:1041-1049.
- Maguire RO and JT Sims. 2002. Soil testing to predict phosphorus leaching. *J Environ Qual* 31: 1601-1609.
- Mehlich A. 1972. Uniformity of expressing soil test results. A case for calculating results on a volume basis. *Commun Soil Sci Plant Anal* 3: 417-424.
- Meetei WH, HS Athokpam, RKK Singh, L Watham, N Chongtham, KN Devi, NB Singh, KL Singh and NJ Singh. 2015. Evaluation of some soil test methods in acid soils for available phosphorus for soybean of Imphal East Distric, Manipur, India. *Afr J Agric Res* 10: 767-771.
- Morel C, N Ziadi and A Messiga. 2014. Modeling of phosphorus dynamics in contrasting agroecosystems using long-term field experiments. *Can J Soil Sci* 94: 377-387.
- Murphy J and JP Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal Chimica Acta* 27: 31-36.
- Nafiu A. 2009. Effects of soil properties on the kinetics of desorption of phosphate from Alfisols by anion-exchange resins. *J Plant Nutr Soil Sci* 172:101-107.
- OelkersEH and E Valsami-Jones. 2008. Phosphate mineral reactivity and global Sustainability. *Elements* 4: 83-87.
- Olsen SR, CV Cole, FS Watanabe and LA Dean. 1954. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. Cir. No. 939 USDA Washington, DC p. 19.
- Pavlatou A and NA Polyzopoulos. 1988. The role of diffusion in the kinetics of phosphate desorption: the relevance of the Elovich equation. *Eur J Soil Sci* 39: 425-436.
- Rouached H, AB Arpat and Y Poirier. 2010. Regulation of phosphate starvation responses in plants: signaling players and cross-talks. *Mol Plant* 3:288-299.
- Schachtman DP, RJ Reid and SM Ayling. 1998. Phosphorus uptake by plants: from soil to cell. *Plant Physiol* 116: 447-453.
- Schoumans OF. 2013. Description of the phosphorus sorption and desorption processes in lowland peaty clay soils. *Soil Sci* 178: 291-300.
- Singh BR, T Krogstad, YS Shivay, BG Shivakumar and M Bakkegard. 2005. Phosphorus fractionation and sorption in P-enriched soils of Norway. *Nutr Cycl Agroecosyst* 73: 245-256.

- Soil Research Institute. 2005. Technical Guidelines for Chemical Analysis of Soil, Plant, Water, and Fertilizer. Soil Research Institute. Bogor. 136 p.
- Sparks DL. 1989. Kinetics of Soil Chemical Processes. Academic Press. INC. 210 p.
- Tiesen H and JO Moir. 2008. Characterization of Available P by Sequential Extraction. In: MR Carter and EG Gregorich (eds). Soil Sampling and Methods of Analysis. 2nd Edition. Canadian Society of Soil Sciences, pp. 293-306
- Thom WO and M Utomo. 1991. Laboratory Management and Soil and Plant Analysis Methods. University of Lampung. 85 pp (in Indonesian).
- Tipping E, S Benham and JF Boyle. 2014. Atmospheric deposition of phosphorus to land and freshwater. *Environ Sci: Processes & Impacts* 16: 1608-1617.
- Torrent J and A Delgado. 2001. Using phosphorus concentration in the soil solution to predict phosphorus desorption to water. *J Environ Qual* 30: 1829-1835.
- Van der Zee SEATM and WH van Riemsdijk. 1986. Sorption kinetics and transport of phosphate in sandy soil. *Geoderma* 38: 293-309.
- Van der Zee SEATM and WH van Riemsdijk. 1988. Model for long-term phosphate reaction kinetics in soil. *J Environ Qual* 17: 35-41.
- Vetterlein D, C Bergmann and RF Hüttl. 1999. Phosphorus availability in different types of open-cast mine spoil and the potential impact of organic matter application. *Plant Soil* 213: 189-194.
- Wahida A, A Fahmi and A Jamberi. 2007. Effect of phosphate rock fertilizer from Morocco on the rice growth in land acid sulfate. *J Trop Soil* 12: 85-90 (in Indonesian).
- Wolf AM and DE Baker. 1985. Comparison of soil test phosphorus by Olsen, Bray-P1, Mehlich-1, and Mehlich-3 methods. *Commun Soil Sci Plant Anal* 16: 467-484.
- Ziadi N, RR Simard, TS Tran and G Allard. 2001. Soil-available phosphorus as evaluated by desorption techniques and chemical extractions. *Can J Soil Sci* 81: 167-174.