

# Effect of Amendments on Soil Charge Characteristics in Sub-Optimal Dryland of Aceh Besar, Indonesia

<sup>1\*</sup>Sufardi Sufardi, <sup>1</sup>Teti Arabia, <sup>1</sup>Khairullah Khairullah, <sup>1</sup>Tri Chairina

<sup>1</sup>Department of Soil Science, Faculty of Agriculture, Universitas Syiah Kuala, Jalan Tgk Hasan Krueng Kalee 3, Darussalam, Banda Aceh 23111

sufardi\_usk@usk.ac.id

## ABSTRACT

Soil colloids play a critical role in controlling chemical behavior, particularly in tropical sub-optimal drylands dominated by variable-charge soils. However, information on the comparative effects of organic and inorganic amendments on soil charge characteristics across different soil orders remains limited. This study aimed to evaluate the effects of compost, biochar, CaCO<sub>3</sub>, and Superphosphate fertilizer (SP-36) on the charge characteristics of tropical dryland soils. Nine amendment combinations were applied to topsoil (0–20 cm) from Andisols (Saree), Inceptisols (Cucum), and Ultisols (Jantho), followed by incubation for 60 days at 25–27 °C under field capacity conditions. Soil charge parameters were analyzed before and after incubation. Results showed that amendment type and rate significantly affected soil charge characteristics, with responses varying among soil orders. Organic amendments (15–30 t ha<sup>-1</sup>) and SP-36 (2–4 t ha<sup>-1</sup>) decreased pH<sub>0</sub> and significantly increased cation exchange capacity (CEC) and net negative charge, indicating improved soil reactivity. In contrast, CaCO<sub>3</sub> increased pH<sub>0</sub> and reduced negative charge. These findings support the strategic use of organic amendments to enhance soil fertility and sustainability.

**Keywords:** Soil amendments, Soil colloids, Surface charge, Tropical drylands, Variable-charge soils

## ABSTRAK

Koloid tanah memainkan peran penting dalam mengendalikan perilaku kimia tanah, terutama pada lahan kering suboptimal tropis yang didominasi oleh tanah bermuatan variabel. Namun, informasi mengenai perbandingan pengaruh bahan amelioran organik dan anorganik terhadap karakteristik muatan tanah pada berbagai ordo tanah masih terbatas. Penelitian ini bertujuan untuk mengevaluasi pengaruh kompos, biochar, CaCO<sub>3</sub>, dan SP-36 terhadap karakteristik muatan tanah pada tanah kering

tropis. Sebanyak sembilan kombinasi amelioran diaplikasikan pada tanah lapisan atas (0–20 cm) dari Andisols (Saree), Inceptisols (Cucum), dan Ultisols (Jantho), kemudian diinkubasi selama 60 hari pada suhu 25–27 °C dalam kondisi kapasitas lapang. Parameter muatan tanah dianalisis sebelum dan setelah inkubasi. Hasil penelitian menunjukkan bahwa jenis dan dosis amelioran berpengaruh nyata terhadap karakteristik muatan tanah, dengan respons yang berbeda antarordo tanah. Amelioran organik (15–30 t ha<sup>-1</sup>) dan SP-36 (2–4 t ha<sup>-1</sup>) menurunkan nilai pH<sub>0</sub> dan secara signifikan meningkatkan kapasitas tukar kation (KTK) serta muatan negatif bersih, yang menunjukkan peningkatan reaktivitas tanah. Sebaliknya, CaCO<sub>3</sub> meningkatkan pH<sub>0</sub> dan menurunkan muatan negatif. Temuan ini mendukung penggunaan amelioran organik secara strategis untuk meningkatkan kesuburan tanah dan ketersediaan hara bagi tanaman.

**Kata kunci:** Amelioran tanah, Koloid tanah, Muatan permukaan, Lahan kering tropis, Tanah bermuatan variabel.

## INTRODUCTION

Suboptimal land (marginal lands) refers to land with inherently low productivity due to various limiting factors that hinder crop production, particularly food crops (Csikós and Tóth, 2023). In Indonesia, suboptimal drylands occur across both dry and humid climatic zones, with the largest proportion located in humid tropical environments, commonly referred to as tropical drylands. These lands represent an important potential resource for agricultural expansion; however, many have undergone significant degradation and are currently classified as critical land, covering more than 18 million hectares (Pitaloka, 2020; Arista et al., 2023). Improving the quality and productivity of these lands has therefore become a key priority for sustainable agricultural development and food security in tropical regions (Samijan et al., 2023; Sufardi, 2024; Sufardi and Helmi, 2025).

In Aceh Besar District, Indonesia, spatial analysis indicates that suboptimal dryland covers approximately 239,387.91 hectares (Umar et al., 2021). Nevertheless, only part of this area can be used for agriculture because large portions are in forests and conservation areas. Previous studies reported that the soils of this region consist of several soil orders, including Ultisols, Oxisols,

Inceptisols, Entisols, and Andisols, with Mollisols occurring in limited areas (Sufardi et al., 2020). These soils are generally characterized by low clay activity due to mineralogical compositions dominated by Fe and Al hydroxide-oxides (Fitri et al., 2020; Cao et al., 2024, Quantin and Becquer, 2025). As a consequence, the soils typically exhibit low negative surface charge, low cation exchange capacity (CEC), and relatively high zero point of charge ( $pH_o$ ) values, which are often higher than the actual soil pH (Kautsar et al., 2018; Sufardi et al., 2019; Sufardi et al., 2020).

Many soils in tropical suboptimal drylands are classified as variable-charge soils. In these soils, the surface charge of soil colloids originates from hydroxyl groups associated with clay minerals, phyllosilicates, allophane–imogolite complexes, hydrous oxides, and soil organic matter (Sparks et al., 2022). Unlike permanent-charge soils, the magnitude and even the sign of the surface charge in variable-charge soils can change depending on soil pH and electrolyte concentration (Alemayehu and Teshome, 2021). These charge characteristics strongly influence soil chemical processes, including nutrient retention, ion exchange, and the adsorption behavior of cations and anions (Khanri et al., 2025).

Suboptimal drylands generally have low soil fertility due to unfavorable physicochemical properties. These soils typically contain low organic matter, limited availability of essential nutrients such as phosphorus and potassium, low CEC, and acidic soil reactions (McLeod et al., 2020; Sufardi et al., 2022; Abdullah et al., 2025; Sufardi et al., 2025). The low fertility status of these soils is closely related to the charge characteristics of soil colloids (Uehara and Gillman, 1985; Yu et al., 2025; Violante and Gianfreda, 2023). In variable-charge soils, the relatively high zero point of charge significantly affects nutrient dynamics, particularly the adsorption of anions such as phosphate on soil colloid surfaces (Nguyen et al., 2020; Penn and Camberato, 2019). Consequently, improving soil charge characteristics is essential for enhancing nutrient availability and soil fertility in tropical dryland ecosystems (Sufardi, 2024; Usman, 2024).

One of the common approaches to improving the physicochemical properties of such soils is the application of soil amendments. Amendments are used to reduce soil acidity, increase negative

surface charge, enhance CEC, and modify the zero point of charge of soil colloids. However, previous studies have shown that different types of amendments may produce contrasting effects on the charge characteristics of variable-charge soils (Wen et al., 2020; Sufardi et al., 2024). For example, lime or  $\text{CaCO}_3$  application to acidic tropical soils can increase soil pH and CEC while reducing exchangeable Al (Guddisa et al., 2018; Xia et al., 2020). Liming with  $\text{CaCO}_3$  is widely used to ameliorate acidic soils by increasing soil pH, neutralizing exchangeable Al, and increasing base saturation. The increase in soil pH promotes the development of pH-dependent negative charges on soil colloids, which subsequently enhances the cation exchange capacity (CEC), particularly in highly weathered soils dominated by variable charge minerals (Makaza et al., 2026; Moreira et al., 2026). On the other hand, liming may also increase the soil  $\text{pH}_0$  value because  $\text{CaCO}_3$  itself has a relatively high zero point of charge (El-Aswad et al., 2023).

Organic amendments are also widely used to improve soil physicochemical properties. Organic materials can increase the density of negative surface charges on soil colloids, thereby enhancing nutrient retention, although their effect on increasing soil pH is generally limited (Li et al., 2021; Lin et al., 2025). Theoretically, the magnitude of variable charge is influenced by several factors, including soil pH, electrolyte concentration, ionic valence, temperature, and the inherent  $\text{pH}_0$  status of the soil (Van Ranst et al., 2017; Patel, 2025). Despite the recognized importance of soil amendments in improving soil quality, comparative information on how different organic and inorganic amendments influence soil charge characteristics across various soil orders in tropical suboptimal drylands remains limited.

Although numerous studies have investigated the role of soil amendments in improving the chemical properties of tropical soils, most of these studies have primarily focused on changes in soil pH, nutrient availability, and crop productivity. Comparatively little attention has been given to how different types of amendments influence the surface charge characteristics of soil colloids, particularly in variable-charge soils that dominate tropical suboptimal drylands. Furthermore, previous studies generally examined individual soil types or single amendments, while comparative

studies across different soil orders within the same agro-ecological region remain limited. This limitation restricts a comprehensive understanding of how organic and inorganic amendments modify the electrochemical behavior of soil colloids and subsequently influence soil fertility. Therefore, investigating the response of soil charge characteristics to different amendments across multiple soil orders is essential for developing more effective soil management strategies in tropical suboptimal drylands.

This study offers new insights into the role of organic and inorganic amendments in regulating soil surface charge properties in tropical dryland soils, which is essential for developing sustainable soil management strategies in suboptimal agricultural lands. Therefore, this study aims to evaluate the effects of several soil amendments on the charge characteristics of soils in the suboptimal drylands of Aceh Besar, Indonesia.

## MATERIALS AND METHODS

The research was conducted under laboratory conditions using topsoil (0–20 cm) collected from three soil orders in sub-optimal dryland areas of Aceh Besar Regency, Aceh, Indonesia. The soils consisted of Andisols (Eutric Hydrudands) from Saree (05°27'03.2" N; 95°43'45.2" E), Inceptisols (Oxyaquic Dystrudepts) from Cucum Village (05°18'18" N; 95°32'48" E), and Ultisols (Typic Kandiuults) from Jantho (05°16'58" N; 95°37'51.8" E). Soil samples were collected from a depth of 0–20 cm. Before treatment application, the samples were air-dried for one week, ground, and sieved through a 0.5-mm mesh. The experiment was arranged in a completely randomized design (CRD) with two factors. The first factor was the type and rate of soil amendment with nine levels: control (without amendment), biochar at 15 t ha<sup>-1</sup>, biochar at 30 t ha<sup>-1</sup>, compost at 15 t ha<sup>-1</sup>, compost at 30 t ha<sup>-1</sup>, CaCO<sub>3</sub> at 2 t ha<sup>-1</sup>, CaCO<sub>3</sub> at 4 t ha<sup>-1</sup>, SP-36 at 2 t ha<sup>-1</sup>, and SP-36 at 4 t ha<sup>-1</sup>. The second factor consisted of three soil orders: Andisols (Saree), Inceptisols (Cucum), and Ultisols (Jantho). These two factors resulted in 27 treatment combinations with a total of 54 experimental units.

For the incubation experiment, 500 g of soil were placed into plastic bottles. Biochar, compost, CaCO<sub>3</sub> (lime), and phosphate fertilizer (SP-36) were then applied according to the designated

treatments and thoroughly mixed with the soil. Distilled water was added to each bottle to adjust the soil moisture to field capacity. The samples were incubated for 60 days in a temperature-controlled room at 25–27 °C. After the incubation period, soil samples were removed from the bottles and air-dried for one week. The dried soils were then gently crushed and sieved through a 1.0-mm mesh. Subsequently, 100 g of soil from each sample was collected for laboratory analysis. To determine the initial soil charge characteristics, soil samples from each soil order were also analyzed before the incubation experiment.

The soil charge characteristics analyzed included  $pH_0$ ,  $pH(H_2O)$ ,  $pH(KCl)$ ,  $\Delta pH(pH KCl - pH H_2O)$ , variable charge, cation exchange capacity (CEC), and exchangeable cations (Ca, Mg, K, Na, Al, and H). The zero point of charge ( $pH_0$ ) was determined using the ion adsorption and potentiometric titration method with  $CaCl_2$ , following the procedure of Uehara and Gillman (1985). Soil pH in  $H_2O$  and KCl was measured using a pH meter. The variable charge was calculated as the difference between the CEC determined by the 1 N  $NH_4OAc$  extraction at pH 7 and the effective CEC, which was obtained from the sum of exchangeable base cations and exchangeable acidity (Al + H) extracted with 1 M KCl.

The data were analyzed using analysis of variance (ANOVA), and treatment means were compared using the least significant difference (LSD) test at  $P < 0.05$  (Gomez and Gomez, 1984).

## RESULTS AND DISCUSSION

### The initial Soil Characteristics

Table 1 shows that the Andisol from Saree is characterized by low pH ( $H_2O$  and KCl) and very low available P, but contains high organic C, moderate total N, and very high potential  $P_2O_5$  and  $K_2O$ . These results indicate that the main constraints of this soil are acidic pH and low available P. In the Inceptisol from Cucum Village, the major limitations include slightly acidic pH, very low available P, low organic C and total N, and low  $K_2O$  content. Similarly, the Ultisol from Jantho exhibits slightly acidic pH and very low levels of available P, organic C, and total N. Based on these chemical properties, the Inceptisol (Cucum) and Ultisol (Jantho) are classified as having low soil fertility status

due to low soil pH, low organic C, and low base saturation, despite their relatively high potential  $P_2O_5$  content. Another characteristic of these sub-optimal dryland soils is the relatively high zero point of charge ( $pH_0$ ), which exceeds the soil pH measured in  $H_2O$ . This condition indicates that the three soil orders used in this study are dominated by variable-charge soils, as reported by Wen et al. (2020).

### **Effects of Soil Amendments on Charge Characteristics**

#### *pH $H_2O$ , pH KCl, and $\Delta pH$*

The incubation experiment demonstrated that the application of different types and rates of amendments significantly influenced soil pH ( $H_2O$  and KCl) in sub-optimal dryland soils of Aceh Besar Regency, Aceh, Indonesia (Table 2). However, the magnitude of pH change varied among soil orders. In the Andisol from Saree, most amendments did not significantly change soil pH. A slight increase in pH was observed only with the application of compost at  $30\text{ t ha}^{-1}$  and  $CaCO_3$  at  $2\text{--}4\text{ t ha}^{-1}$ . The limited change in pH may be related to the high buffering capacity of Andisols, which are dominated by amorphous minerals and organic matter that tend to stabilize soil acidity (Parada et al., 2024; Fajarini et al., 2025). Without amendment, the soil pH was acidic (5.41), while the addition of compost and  $CaCO_3$  increased the pH slightly to 5.82, which is still classified as slightly acidic. The liming effect of  $CaCO_3$  occurs through the neutralization of exchangeable acidity and the precipitation of  $Al^{3+}$ , thereby increasing soil pH (Zhang et al., 2023; Abdi, 2024).

In the Inceptisol from Cucum Village, an increase in soil pH was observed only with  $CaCO_3$  application at rates of  $2\text{--}4\text{ t ha}^{-1}$ , indicating that liming was the most effective treatment for reducing soil acidity. In contrast, compost and biochar did not significantly affect soil pH. The application of SP-36 at  $2\text{--}4\text{ t ha}^{-1}$  tended to decrease soil pH from slightly acidic to acidic conditions, which may be associated with the release of acidity during phosphate fertilizer reactions in the soil.

Similarly, in the Ultisol from Jantho, soil pH increased with the application of compost and  $CaCO_3$ , whereas biochar had no significant effect. The increase in pH with compost application may be related to the release of basic cations during organic matter decomposition, which can partially neutralize soil acidity. Conversely, the application of SP-36 tended to reduce soil pH, indicating its potential

acidifying effect in highly weathered soils. These results suggest that liming and organic amendments are more effective in improving soil acidity in variable-charge soils compared with biochar or phosphate fertilizer alone.

The incubation experiment indicated that lime ( $\text{CaCO}_3$ ) and compost amendments were effective in increasing soil pH ( $\text{H}_2\text{O}$ ), whereas biochar and phosphate fertilizer (SP-36) were less effective in improving soil acidity in sub-optimal dryland soils of Aceh Besar Regency, Aceh, Indonesia. Compost has a high potential to restore soil fertility because it serves as an important source of soil organic matter. During incubation, the decomposition of organic materials releases basic cations and organic ligands that can neutralize acidic cations such as  $\text{H}^+$  and  $\text{Al}^{3+}$ , thereby increasing soil pH, as reported by Zhang et al. (2023). Similarly,  $\text{CaCO}_3$  application increased soil pH through the dissociation of  $\text{CaCO}_3$  into  $\text{Ca}^{2+}$  and  $\text{CO}_3^{2-}$  ions. The carbonate ions react with  $\text{H}^+$  and  $\text{Al}^{3+}$  in the soil solution, leading to the precipitation of Al compounds and a reduction in exchangeable acidity, which ultimately increases soil pH (Panda et al., 2025; Makaza et al., 2026).

In contrast, the application of biochar at rates of 15–30  $\text{t ha}^{-1}$  did not significantly increase soil pH in this study. This may be related to the characteristics of the biochar used, which contains a high proportion of stable carbon and functions primarily as an adsorbent rather than as a liming material, as suggested by Tang (2025). However, other studies have reported that biochar can increase soil pH depending on the application rate and feedstock characteristics (Juriga et al., 2022). Biochar may contribute to soil pH neutralization and improve soil chemical properties, particularly in acidic soils, while also serving as a stable carbon source due to its resistance to microbial decomposition (Yang et al., 2025).

The application of SP-36 at rates of 2–4  $\text{t ha}^{-1}$  tended to decrease soil pH. This effect may be associated with the presence of sulfate in the fertilizer and the formation of phosphoric acid during phosphate reactions in the soil solution. Nevertheless, other studies have shown that phosphate application can also increase soil pH under certain conditions (Zhao et al., 2021). This occurs when

phosphate anions are adsorbed onto positively charged soil colloids, which reduces proton activity and consequently increases soil pH (Zhang et al., 2023; Sufardi et al., 2024).

Table 2 also shows that the types and rates of amendments affected soil pH (KCl), although the magnitude of this effect varied among soil orders. In the Andisol from Saree, the application of compost at 30 t ha<sup>-1</sup> increased pH (KCl) compared with the control treatment without amendment. In the Inceptisol from Cucum Village, the application of CaCO<sub>3</sub> at rates of 2–4 t ha<sup>-1</sup> increased soil pH (KCl) relative to the control. Similar findings were reported by Ameyu (2019), who showed that the application of CaCO<sub>3</sub> combined with humic substances increased soil pH to above 6.5 and reduced exchangeable Al due to precipitation with CaCO<sub>3</sub> and chelation by functional groups of humic compounds.

The increase in soil pH following CaCO<sub>3</sub> application is mainly attributed to the neutralization of soil acidity through the replacement of H<sup>+</sup> ions by Ca<sup>2+</sup> on the exchange complex, as reported by Mosharrof et al. (2021). In the Ultisol from Jantho, the application of CaCO<sub>3</sub> at 4 t ha<sup>-1</sup> also increased soil pH (KCl) compared with the control. Liming can produce hydroxyl ions (OH<sup>-</sup>) in the soil solution, which reduce exchangeable Al through the formation of insoluble Al(OH)<sub>3</sub> precipitates, thereby decreasing soil acidity (Guddisa et al., 2018; Panda et al., 2025; Makaza et al., 2026).

The application of amendments to variable-charge soils in the sub-optimal drylands of Aceh Besar Regency also influenced changes in ΔpH (pH KCl – pH H<sub>2</sub>O), although the effect was not significant in the Andisol from Saree (Table 2). In the Inceptisol from Cucum, the application of biochar, compost, CaCO<sub>3</sub>, and phosphate fertilizer (SP-36) increased ΔpH values compared with the control treatment. In contrast, in the Ultisol from Jantho, only biochar and compost increased ΔpH values.

An increase in ΔpH indicates a greater development of negative charge on soil colloids (Wen et al., 2020), which can improve soil chemical properties and nutrient retention. The increase in negative surface charge following amendment application depends not only on the type and rate of amendment but also on soil mineralogical composition, which differs among soil orders (Sufardi et al., 2019; Belghazdis and Hachem, 2022).

The changes in soil pH following amendment application were closely related to variations in  $\Delta\text{pH}$ , cation exchange capacity (CEC), and the development of variable surface charge. In variable-charge soils, the magnitude and sign of surface charge depend strongly on soil pH relative to the zero point of charge ( $\text{pH}_0$ ). When soil pH exceeds  $\text{pH}_0$ , negative charges on soil colloids increase, leading to greater CEC and enhanced nutrient retention. Conversely, when soil pH is lower than  $\text{pH}_0$ , positive charges tend to dominate the colloidal surfaces. The increase in soil pH observed after the application of compost and  $\text{CaCO}_3$  likely promoted the development of negative surface charges, as indicated by the increase in  $\Delta\text{pH}$  values. This condition enhances the dissociation of functional groups on soil organic matter and variable-charge minerals such as Fe and Al oxides, thereby increasing the number of negatively charged sites available for cation adsorption. As a result, the soil's capacity to retain essential nutrient cations also increases.

The application of biochar, compost, and phosphate fertilizer (SP-36) tended to increase soil pH, which consequently increased  $\Delta\text{pH}$  values ( $\text{pH}_{\text{KCl}} - \text{pH}_{\text{H}_2\text{O}}$ ) and promoted the development of more negative surface charges on soil colloids. However, the effect of  $\text{CaCO}_3$  amendment on soil charge was less consistent. At higher application rates,  $\text{CaCO}_3$  may increase the concentration of  $\text{Ca}^{2+}$  in the soil solution, which can enhance cation adsorption and partially increase the positive charge on soil colloid surfaces.

The development of negative charge on soil colloids plays an important role in maintaining soil fertility because negatively charged sites can adsorb and retain nutrient cations, thereby reducing nutrient losses through leaching (Mosharrof et al., 2021). This phenomenon is commonly observed in highly weathered soils with variable-charge characteristics in tropical and subtropical regions, where soil surface charge is strongly influenced by soil pH and mineral composition (Van Ranst et al., 2017; Cao et al., 2024).

Furthermore, the relationship between soil pH and the zero point of charge ( $\text{pH}_0$ ) is an important indicator of soil charge behavior. The difference between  $\text{pH}_0$  and soil pH ( $\text{pH}_0 - \text{pH}_{\text{H}_2\text{O}}$ ) reflects the dominance of positive or negative charges on soil colloids. When soil pH approaches or exceeds

$pH_0$ , negative charges tend to dominate, whereas when soil pH is lower than  $pH_0$ , positive charges become more dominant. Therefore, amendments that increase soil pH can shift the balance toward greater negative surface charge in variable-charge soils.

Overall, the results demonstrate that soil amendments play an important role in modifying the surface charge characteristics of variable-charge soils. The increase in soil pH following the application of compost and  $CaCO_3$  shifted the balance between positive and negative charges on soil colloids, as reflected by changes in  $\Delta pH$  and the relationship between soil pH and the zero point of charge ( $pH_0$ ). When soil pH approaches or exceeds  $pH_0$ , the number of negative surface charges increases due to the dissociation of functional groups on soil organic matter and oxide minerals, which consequently enhances cation exchange capacity (CEC) and nutrient retention.

#### *$pH_0$ , and $pH_0 - pH_{H_2O}$*

The effects of amendment materials on the charge characteristics of sub-optimal dryland soils in Aceh Besar, as indicated by the parameters  $pH_{H_2O}$ ,  $pH_0$ , and the difference between  $pH_0$  and  $pH_{H_2O}$ , are presented in Figures 1 and 2. Figure 1 shows that amendments of different types and application rates influenced the  $pH_0$  of variable-charge soils, with the magnitude of the effect depending on both amendment type and soil order. In general, the application of amendments is expected to reduce soil  $pH_0$ . The results of this experiment indicate that organic amendments, namely biochar and compost, as well as phosphate fertilizer (SP-36), were effective in decreasing soil  $pH_0$ , whereas  $CaCO_3$  application increased soil  $pH_0$ . In Andisols from Saree, the application of biochar, compost, and SP-36 at specific rates reduced  $pH_0$  from 4.01 to 3.06, and similar trends were observed in Inceptisols from Cucum and Ultisols from Jantho. In contrast,  $CaCO_3$  application increased  $pH_0$  from 4.05 to 6.00. These results demonstrate that organic and phosphate amendments can improve the charge characteristics of soil colloids by reducing soil  $pH_0$ . When the actual soil pH is lower than  $pH_0$ , soil colloids are predominantly positively charged (Wen et al., 2020; Yu et al., 2025).

Conversely, when soil pH exceeds  $pH_0$ , soil colloids become negatively charged (Lal, 2017). Most soils in humid tropical regions generally have relatively high  $pH_0$  values. Consequently, anions in the

soil solution, such as phosphate, molybdate, and sulfate, tend to be strongly adsorbed onto the surfaces of soil particles, thereby reducing their availability in the soil solution (Barrow and Hartemink, 2023). The increase in soil  $pH_0$  following lime application can be attributed to the carbonate anions derived from  $\text{CaCO}_3$ , which inherently possess a high  $pH_0$  and thus contribute to increasing the soil  $pH_0$  when applied (Guddisa et al., 2018). Furthermore,  $\text{Ca}^{2+}$  cations released from lime may be adsorbed onto the inner layer of the colloidal surface, altering the surface charge of soil colloids and increasing the point of zero charge.

The results of this experiment also showed that a decrease in soil  $pH_0$  following amendment application, particularly with organic materials, was accompanied by an increase in soil negative charge. As shown in Figure 2, except for the Andisols from Saree, organic amendments (biochar and compost) applied at rates of 10–20  $\text{t ha}^{-1}$  increased the  $pH_0 - \text{pH H}_2\text{O}$  value, indicating an increase in the net negative charge of the soil. Among the tested amendments, compost produced the greatest increase in negative charge, followed by biochar and phosphate fertilizer. The highest negative charge, expressed as the difference between  $pH_0$  and  $\text{pH H}_2\text{O}$ , was obtained with the application of compost at 20  $\text{t ha}^{-1}$ . Under this treatment, the negative charge increased from  $-1.00$  to  $-4.00$ , representing a fourfold increase. This increase in negative charge is associated with the inherently low  $pH_0$  of organic matter and phosphate compounds (Koishi et al., 2020). Organic matter contains humic and fulvic substances that contribute to the neutralization of positive charges on soil colloids (Li et al., 2021). In contrast, although  $\text{CaCO}_3$  application increases soil pH, it simultaneously increases soil  $pH_0$ , resulting in a tendency toward a more positive net colloidal charge, particularly at higher application rates, as indicated by the relatively small difference between  $pH_0$  and  $\text{pH H}_2\text{O}$  (Tang et al., 2025; Zhang et al., 2025).

Overall, these findings highlight that organic amendments, particularly compost and biochar, play a crucial role in modifying the charge characteristics of variable-charge soils by reducing  $pH_0$  and increasing net negative surface charge, thereby improving nutrient retention in sub-optimal tropical drylands.

### *CEC and ECEC*

This experiment also demonstrated that amendment application influenced both the cation exchange capacity (CEC) and effective cation exchange capacity (ECEC) of the soils. As shown in Figure 3, different types and application rates of amendments affected CEC, although the magnitude of the effect varied among amendment types and soil orders. The application of biochar, compost, and phosphate fertilizer increased potential CEC, particularly in the Inceptisols of Cucum and Ultisols of Jantho. In contrast, the effect was relatively small in the Andisols of Saree, which inherently possess high CEC and strong buffering capacity due to the presence of allophane minerals (Sufardi et al., 2020; Belghazdis and Hachem, 2022). Conversely, the application of  $\text{CaCO}_3$  at rates of 2–4 t ha<sup>-1</sup> tended to reduce soil CEC. The increase in CEC following amendment application is associated with the increase in negative surface charge of soil colloids, as soil CEC is positively correlated with the magnitude of colloidal charge (Khadim et al., 2023). Compost, in particular, contributes additional negative charges derived from carboxyl ( $\text{COO}^-$ ) and hydroxyl ( $\text{OH}^-$ ) functional groups, which enhance the soil's ability to retain cations in the soil solution (Senesi and Loffredo, 2018; Boukhatem et al., 2025). The improvement of CEC in Inceptisols of Cucum and Ultisols of Jantho is particularly important because these soils contain relatively low organic matter. In addition,  $\text{CaCO}_3$  application can increase the concentration of  $\text{Ca}^{2+}$  ions in the soil solution, which occupy exchange sites on the soil exchange complex and help neutralize excess  $\text{Al}^{3+}$  and  $\text{H}^+$  ions (Iqbal, 2012). Overall, these results indicate that organic amendments are more effective than lime in enhancing soil CEC through the increase of negative surface charge in variable-charge soils.

Effective cation exchange capacity (ECEC) also increased following the incubation experiment after the application of amendment materials (Figure 4). ECEC represents the sum of exchangeable base cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{Na}^+$ ) and acidic cations ( $\text{Al}^{3+}$  and  $\text{H}^+$ ) present on the soil exchange complex. The results showed that the three soil orders—Andisols from Saree, Inceptisols from Cucum, and Ultisols from Jantho—exhibited the highest ECEC values when compost was applied at

a rate of 30 t ha<sup>-1</sup>, whereas the lowest ECEC values were observed in the control treatment without amendments. The ECEC values of the studied soils remained relatively low, indicating that the sub-optimal dryland soils of Aceh Besar are classified as variable-charge soils with low-activity clay (LAC), characterized by ECEC values of less than 12 cmol kg<sup>-1</sup>. As shown in Figure 4, the ECEC values ranged from 5.91–7.05 cmol kg<sup>-1</sup> in the Andisols of Saree, 5.08–6.47 cmol kg<sup>-1</sup> in the Inceptisols of Cucum, and 1.65–4.28 cmol kg<sup>-1</sup> in the Ultisols of Jantho.

Soils dominated by low-activity clays are typically composed of kaolinite and hydroxides of Al and Fe, which commonly exhibit positive surface charges under acidic soil conditions and consequently result in relatively low CEC values (Li et al., 2021). In such soils, the negative charge contributing to soil CEC mainly originates from the dissociation of hydrogen ions from hydroxyl (OH) groups associated with Si and Al oxides, as well as from a small proportion of broken crystal edges. However, the charge generated from these processes contributes only a limited amount to the overall soil CEC (Ameyu, 2019; Adetunji and Blanco-Canqui, 2026). Overall, these results indicate that the low ECEC observed in the studied soils is closely related to the dominance of low-activity clay minerals typical of highly weathered tropical soils.

Overall, the results demonstrate that organic amendments, particularly compost and biochar, enhance the electrochemical properties of variable-charge soils by increasing negative surface charge and improving CEC and ECEC. This improvement provides an effective strategy for enhancing nutrient retention and fertility in sub-optimal tropical dryland soils.

## CONCLUSIONS

The results demonstrate that organic amendments significantly improve the charge characteristics and physicochemical properties of variable-charge soils in the sub-optimal drylands of Aceh Besar, including Andisols, Inceptisols, and Ultisols. Application of compost and biochar (15–30 t ha<sup>-1</sup>) combined with phosphate fertilizer (SP-36) enhanced soil pH, increased effective cation exchange capacity (ECEC), and promoted greater development of negative surface charge. These amendments

also reduced soil pH<sub>0</sub>, indicating improved charge balance in variable-charge systems. In contrast, the application of CaCO<sub>3</sub> (2–4 t ha<sup>-1</sup>) increased soil pH and CEC but tended to raise pH<sub>0</sub> and reduce negative surface charge. Overall, organic amendments were more effective than CaCO<sub>3</sub> in improving the electrochemical properties of tropical variable-charge soils, suggesting their important role in enhancing soil fertility and sustainable management of sub-optimal drylands.

## ACKNOWLEDGEMENTS

The authors sincerely acknowledge the support of colleagues and laboratory staff who contributed to field sampling and laboratory analyses. Their technical assistance and constructive support were essential for the completion of this research.

## REFERENCES

- Abdi BT 2024. Studies on the effects of liming acidic soil on improving soil physicochemical properties and yield of crops: A review. *Middle East Research Journal of Agriculture and Food Science*, 4(03), 95-103. <https://doi.org/10.36348/merjafs.2024.v04i03.001>
- Abdullah UH, S Sufardi, Endiyani, R Wilis, Elviani, N Farida, H Akbar 2025. The analysis of plant biomass potential and vegetation diversity in various types of dryland uses in Aceh Besar Regency. *AIP Conference Proceedings* 3277(1), 020004. <https://doi.org/10.1063/5.0269225>
- Adetunji AT, H Blanco-Canqui 2026. Soil health response to biochar combined with other amendments: a review. *Biochar* 8(1), 23. <https://doi.org/10.1007/s42773-025-00531-6>
- Alemayehu B, H Teshome 2021. Soil colloids, types and their properties: A review. *Open Journal of Bioinformatics and Biostatistics* 5(1), 008–013. <https://doi.org/10.17352/ojbb.000010>
- Ameyu T 2019. A review of the potential effect of lime on soil properties and crop productivity improvements. *Journal of Environment and Earth Science* 9(2), 17-23. <https://doi.org/10.7176/jees/9-2-03>
- Arista NID, AD Alfia, H Mubarak, IMSD Arta, DN Rizva, AI Wicaksono 2023. Availability and potential for expansion of agricultural land in Indonesia. *Journal of Sustainability, Society, and Eco-Welfare* 1(1). <https://doi.org/10.61511/jssew.v1i1.2023.242>
- Barrow, N.J. and Hartemink, A.E., 2023. The effects of pH on nutrient availability depend on both soils and plants. *Plant and Soil*, 487(1), 21-37. <https://doi.org/10.1007/s11104-023-05960-5>
- Belghazdis M, EK Hachem 2022. Clay and clay minerals: a detailed review. *International Journal of Recent Technology and Applied Science (IOERTAS)*, 4(2), 54–75. <https://doi.org/10.36079/lamintang.ijortas-0402.367>
- Boukhatem A, O Rached, A Bentellis, S Vasileiadis, P Castaldi, G Garau, S Diquattro 2025. Promoting the recovery of soil health in polluted soils using biochar–compost amendments. *Environmental Science and Pollution Research* 32, 559–574. <https://doi.org/10.1007/s11356-024-35650-3>

- Cao H, X Liu, B Feng, J Sun, D Ma, X Chen, H Li 2024. Effect of pH on the surface charges of permanently/variably charged soils and clay minerals. *Scientific Reports* 14(1), 23169. <https://doi.org/10.21203/rs.3.rs-4684462/v1>
- Csikós N, G Tóth 2023. Concepts of agricultural marginal lands and their utilisation: A review. *Agricultural Systems*, 204, p.103560. <https://doi.org/10.1016/j.agsy.2022.103560>
- El-Aswad AF, MR Fouad, ME Badawy, MI Aly 2023. Effect of calcium carbonate content on potential pesticide adsorption and desorption in calcareous soil. *Communications in Soil Science and Plant Analysis*, 54(10), 1379-1387. <https://doi.org/10.1080/00103624.2022.2146131>
- Fajarini PHN, E Hanudin, BH Purwanto 2025. Analysis of factors affecting differences in potassium buffering capacity in several soil orders. *Agricultural Science/Ilmu Pertanian*, 10(2), 101–112. <https://doi.org/10.22146/ipas.108285>
- Fitri F, T Arabia, S Sufardi 2020. Identification of amorphous Fe, Al, and Si fractions in several soil orders in Aceh Besar dry land with ammonium oxalate extraction. *Jurnal Ilmiah Mahasiswa Pertanian* 4(4), 618–628. <https://doi.org/10.17969/jimfp.v4i4.12689>
- Gomez KA, AAGomez 1984. *Statistical Procedures for Agricultural Research*. John Wiley & Sons.
- Guddisa D, T Tana, A Legesse 2018. Application of different lime rates and phosphorus on soil physicochemical properties of acid soils in western Ethiopia. *Science, Technology and Arts Research Journal* 5(1), 84. <https://doi.org/10.4314/star.v5i1.13>
- Iqbal M 2012. Effect of Al compounds on soil pH and bioavailability of Al in two acid soils. *Turkish Journal of Agriculture and Forestry* 36(6), 720–728. <https://doi.org/10.3906/tar-1109-26>
- Juriga M, E Aydın, J Horák, V Šimanský 2022. Relationships between soil chemical properties and soil structure after initial application and reapplication of biochar and its combination with N fertilization. *Communications in Soil Science and Plant Analysis* 53(1), 114–128. <https://doi.org/10.1080/00103624.2021.1984510>
- Kautsar I, S Ilyas, S Sufardi 2018. Load characteristics and soil physicochemical properties of Ultisols and Andisols in Aceh Besar dry land. *Jurnal Ilmiah Mahasiswa Pertanian* 3(2), 409–419. <https://doi.org/10.17969/jimfp.v3i2.7498>
- Khadim MD, AB Wesal, AW Peezhand 2024. Overview of the impact of compost on bulk density, aggregate consistency, and cation exchange capacity of soils and its consequential effect on crop productivity. *Cognizance Journal of Multidisciplinary Studies*, 4(6), 344-359. <https://doi.org/10.47760/cognizance.2024.v04i06.021>
- Khanri NK, NPN Asri, HA Hendra, S Susantya, DH Susetyo, KH Kirana, E Agustine 2025. Physicochemical characterization of soils in different land uses: A case study in Jatiroke Village, Sumedang, West Java. *Jurnal Penelitian Pendidikan IPA* 11(10), 1116–1131. <https://doi.org/10.29303/jppipa.v11i10.12863>
- Koishi A, L Bragazza, A Maltas, T Guillaume, S Sinaj 2020. Long-term effects of organic amendments on soil organic matter quantity and quality in conventional cropping systems in Switzerland. *Agronomy* 10(12), 1977. <https://doi.org/10.3390/agronomy10121977>
- Lal R 2017. Variable charge soils: Mineralogy and chemistry. In: *Encyclopedia of Soil Science*, Third Edition. CRC Press, pp. 2432–2439. <https://doi.org/10.1081/e-ess3-120053756>
- Li Y, X Dong, EB da Silva, LM de Oliveira, Y Chen, LQ Ma 2017. Mechanisms of metal sorption by biochars: Biochar characteristics and modifications. *Chemosphere*. 178, 466-478. <https://doi.org/10.1016/j.chemosphere.2017.03.072>

- Li Y, Y Su, W Ahmed, S Ren, T Javed, F Yao, X Li 2021. Effects of different organic fertilizers on improving soil from newly reclaimed land to crop soil. *Agriculture* 11(6), 560. <https://doi.org/10.3390/agriculture11060560>
- Lin N, R Jiao, X Li, S Lin, L Wei, B Wu 2025. Surface-engineered biochars enhance soil fertility by modulating microbial assembly and ecological network stability. *Journal of Environmental Management* 395, 127776. <https://doi.org/10.1016/j.jenvman.2025.127776>
- Makaza W, L Khiari, M El Achaby 2026. The meta-analysis study on the effects of the quality of lime materials on the soil physicochemical properties and crop yields in acid soils. *Frontiers in Soil Science* 6, 1725559. <https://doi.org/10.3389/fsoil.2026.1725559>
- McLeod M, S Sufardi, S Harden 2020. Soil fertility constraints and management to increase crop yields in the dryland farming systems of Aceh, Indonesia. *Soil Research* 59(1), 68–82. <https://doi.org/10.1071/SR19324>
- Moreira SG, JRF Gaudencio, FA de Moraes, EG de Moraes, DS Peixoto, HC de Resende, OLV Campos 2026. A practical method for estimating liming requirements based on soil chemical attributes and limestone composition. *Soil and Tillage Research* 255, 106816. <https://doi.org/10.1016/j.still.2025.106816>
- Mosharrof M, MK Uddin, S Jusop, MF Sulaiman, SM Shamsuzzaman, ANA Haque 2021. Changes in acidic soil chemical properties and carbon dioxide emission due to biochar and lime treatments. *Agriculture* 11(3), 219. <https://doi.org/10.3390/agriculture11030219>
- Nguyen TT, QT Dinh, NM Truong 2020. Surface charge characteristics of a variable charge soil (Rhodic Ferralsols) in the Central Highlands. *Vietnam Journal of Science, Technology and Engineering* 62(2), 65–70. [https://doi.org/10.31276/vjste.62\(2\).65-70](https://doi.org/10.31276/vjste.62(2).65-70)
- Panda D, G Sahu, B Bhuyan, S Mishra 2025. Effect of lime on different forms of soil acidity and phosphorus availability in acid soil. *Indian Journal of Ecology* 52(2), 328–333. <https://doi.org/10.55362/ije/2025/4496>
- Parada J, A Neaman, D Zamorano, F Najera, F Matus 2024. Management and liming-induced changes in organo-Al/Fe complexes and amorphous mineral-associated organic carbon: Implications for carbon sequestration in volcanic soils. *Soil and Tillage Research*, 244, 106133. <https://doi.org/10.1016/j.still.2024.106133>
- Patel KK 2025. *Fundamentals of Soil Physical Chemistry*. AG Publishing House.
- Penn CJ, JJ Camberato 2019. A critical review of soil chemical processes that control how soil pH affects phosphorus availability to plants. *Agriculture* 9(6), 120. <https://doi.org/10.3390/agriculture9060120>
- Pitaloka D 2020. Lahan kering dan pola penanaman untuk melestarikan alam. *Jurnal Teknologi Terapan: G-Tech* 2(1), 119–126. <https://doi.org/10.33379/gtech.v2i1.329>
- Quantin C, T Becquer 2025. Tropical soils and sustainable management. In: *Agricultural Soil Science: Sustainable Management of Agricultural Soils*. pp. 249. <https://doi.org/10.1002/9781394361809.ch9>
- Samijan S, Minarsih, S Jauhari, S Basuki, A Susila, E Nurwahyuni, VE Aristya 2023. Revitalizing sub-optimal drylands: Exploring the role of biofertilizers. *Open Agriculture* 8(1), 20220214. <https://doi.org/10.1515/opag-2022-0214>
- Senesi N, E Loffredo 2018. The chemistry of soil organic matter. In *Soil Physical Chemistry* (pp. 239-370). CRC Press.
- Sparks DL, B Singh, MG Siebecker 2022. *Environmental Soil Chemistry*. Elsevier.

- Sufardi S 2024. How to enhance soil quality in dryland farming systems in Indonesia. *IOP Conference Series: Earth and Environmental Science* 1297, 012071. <https://doi.org/10.18805/ijare.af-881>
- Sufardi S, H Helmi 2025. Soil quality index and water stress vulnerability in different land use against drought disaster of Aceh Besar, Indonesia. *Applied and Environmental Soil Science* 2025 (1), 7530709. <https://doi.org/10.1088/1755-1315/1469/1/012030>
- Sufardi S, RB Yakob, M Khalil, T Arabia, K Khairullah 2024. Amelioration's effects on soil chemical properties in maize cultivation in dryland Aceh, Indonesia. *Indian Journal of Agricultural Research* 58(5). <https://doi.org/10.18805/ijare.af-881>
- Sufardi S, S Syafruddin, T Arabia, K Khairullah, HA Umar 2022. Comparison of carbon content in soil and biomass in various types of sub-optimal dryland use in Aceh Besar, Indonesia. *IOP Conference Series: Earth and Environmental Science* 1116, 012049. <https://doi.org/10.1088/1755-1315/1116/1/012049>
- Sufardi S, T Arabia, K Khairullah, K Karnilawati, F Fuadi 2020. Distribution of carbon and soil quality in drylands of Aceh Besar, Indonesia. *IOP Conference Series: Earth and Environmental Science* 458, 012040. <https://doi.org/10.1088/1755-1315/458/1/012040>
- Sufardi S, T Arabia, K Khairullah, K Karnilawati, N Nurnikmat 2019. Distribution of Al, Fe, and Si oxides in three soil orders in the dryland of Aceh Besar, Indonesia. *IOP Conference Series: Earth and Environmental Science* 393, 012081. <https://doi.org/10.1088/1755-1315/393/1/012081>
- Sufardi S, T Arabia, K Khairullah, K Karnilawati, S Sahbudin, Z Zainabun 2020. Charge characteristics and cation exchange properties of hilly dryland soils, Aceh Besar, Indonesia. *Aceh International Journal of Science and Technology* 9(2), 90–101. <https://doi.org/10.13170/aijst.9.2.17565>
- Sufardi S, T Arabia, M Manfarizah, Z Zuraida, Z Zainabun, M Khalil, UH Abdullah 2025. Distribution of soil carbon and nitrogen under primary and cultivated forests in Aceh Besar, Indonesia. *IOP Conference Series: Earth and Environmental Science* 1469, 012030. <https://doi.org/10.1088/1755-1315/1469/1/012030>
- Tang KHD 2025. Biochar amendments for soil restoration: Impacts on nutrient dynamics and microbial activity. *Environments* 12(11), 425. <https://doi.org/10.3390/environments12110425>
- Uehara G, G Gillman 1985. The mineralogy, chemistry, and physics of tropical soils with variable charge clays. *Soil Science* 139(4), 380. <https://doi.org/10.1097/00010694-198504000-00019>
- Umar U, S Sufardi, S Syafruddin, T Arabia 2021. Mapping distribution approach in various types of use of sub-optimal dryland in Aceh Besar District. *IOP Conference Series: Earth and Environmental Science* 715, 012026. <https://doi.org/10.1088/1755-1315/715/1/012026>
- Usman S 2024. Soil and water management perspectives for tropical and dryland areas of Africa. *Soil Studies* 13(2), 103–117. <https://doi.org/10.21657/soilst.1601786>
- Van Ranst E, NP Qafoku, A Noble, RK Xu 2017. Variable charge soils: mineralogy and chemistry. In *Encyclopedia of soil science* (pp. 2432-2439). CRC Press Taylor & Francis. [https://doi.org/10.1016/S0065-2113\(04\)84004-5](https://doi.org/10.1016/S0065-2113(04)84004-5)
- Violante A, L Gianfreda 2023. Adsorption of phosphate on variable charge minerals: competitive effect of organic ligands. In: *Environmental Impacts of Soil Component Interactions*. CRC Press, pp. 29–38. <https://doi.org/10.1201/9780138757045-4>
- Walker DJ, R Clemente, MP Bernal 2004. Contrasting effects of manure and compost on soil pH, heavy metal availability, and growth of *Chenopodium album* L. in a soil contaminated by pyritic mine waste. *Chemosphere* 57(3), 215–224. <https://doi.org/10.1016/j.chemosphere.2004.05.020>

- Wen Y, C Wang, W Cheng, Z Liu, Q Ma, X Zhang, H Tao 2020. Surface charge properties of variable charge soils influenced by environmental factors. *Applied Clay Science* 189, 105522. <https://doi.org/10.1016/j.clay.2020.105522>
- Xia M, M Riaz, M Zhang, B Liu, Z El-Desouki, C Jiang 2020. Biochar increases the nitrogen use efficiency of maize by relieving aluminum toxicity and improving soil quality in acidic soil. *Ecotoxicology and Environmental Safety* 196, 110531 <https://doi.org/10.1016/j.ecoenv.2020.110531>
- Yang W, L Zhang, Z Wang, J Zhang, P Li, L Su 2025. Effects of biochar and nitrogen fertilizer on microbial communities, CO<sub>2</sub> emissions, and organic carbon content in soil. *Scientific Reports*, 15(1), p.9789.
- Yu L, A He, R Tian, X Liu, H Li 2025. Quantitatively distinguishing colloidal composition, surface property of differently sized particles for permanently and variably charged soils. *Applied Clay Science* 265, 107651. <https://doi.org/10.1016/j.clay.2024.107651>
- Zhang L, L Chang, H Liu 2025. Biochar application to soils can regulate soil phosphorus availability: A review. *Biochar* 7, 13. <https://doi.org/10.1007/s42773-024-00415-1>
- Zhang S, Q Zhu, W de Vries, GH Ros, X Chen, MA Muneer, F Zhang, L Wu 2023. Effects of soil amendments on soil acidity and crop yields in acidic soils: A worldwide meta-analysis. *Journal of Environmental Management* 345, 118531. <https://doi.org/10.1016/j.jenvman.2023.118531>
- Zhao H, X Li, G Huang, W Sun, X Qin, S Wei, Z Ye 2021. Effects of various phosphorus fertilizers on maize yield and phosphorus uptake in soils with different pH values. *Archives of Agronomy and Soil Science*. 68(12), 1746-1754 <https://doi.org/10.1080/03650340.2021.1926997>

Accepted Manuscript (May 2025)

## TABLES, FIGURES, AND ILLUSTRATIONS

Table 1. Chemical characteristics of the topsoil of Andisol, Inceptisol, and Ultisol before the experiment

Soil chemical characteristics	Andisols Saree	Inceptisols Cucum	Ultisols Jantho
pH H <sub>2</sub> O (1:2.5)	5.56 <sup>A</sup>	6.45 <sup>SA</sup>	5.97 <sup>SA</sup>
pH KCl (1:2.5)	4.35 <sup>HA</sup>	4.73 <sup>A</sup>	4.01 <sup>HA</sup>
ZPC = pH <sub>o</sub>	5.91 <sup>H</sup>	4.48 <sup>M</sup>	5.58 <sup>H</sup>
CEC (cmol kg <sup>-1</sup> )	31.60 <sup>H</sup>	21.20 <sup>M</sup>	20.40 <sup>M</sup>
Available P (mg kg <sup>-1</sup> )	2.30 <sup>VL</sup>	0.50 <sup>VL</sup>	2.85 <sup>VL</sup>
Organic C (g kg <sup>-1</sup> )	39.5 <sup>H</sup>	0.64 <sup>VL</sup>	0.53 <sup>VL</sup>
Total N (g kg <sup>-1</sup> )	2.52 <sup>M</sup>	0.51 <sup>VL</sup>	0.45 <sup>VL</sup>
Total P <sub>2</sub> O <sub>5</sub> (mg 100 g <sup>-1</sup> )	83.12 <sup>VH</sup>	77.35 <sup>VH</sup>	221.21 <sup>VH</sup>
Total K <sub>2</sub> O (mg 100 g <sup>-1</sup> )	65.98 <sup>VH</sup>	38.25 <sup>L</sup>	28.14 <sup>L</sup>
Base saturation (%)	43.52 <sup>H</sup>	36.74 <sup>M</sup>	25.78 <sup>L</sup>

HA (highly acid), A (acid), SA (slightly acid), VL (very low), M (medium), H (high), VH (very high)

Table 2. pH H<sub>2</sub>O, pH KCl, and ΔpH of three soil orders on various amendment treatments in Aceh Besar dryland

Amendments application	Andisols Saree			Inceptisols Cucum			Ultisols Jantho		
	pH H <sub>2</sub> O	pH KCl	ΔpH	pH H <sub>2</sub> O	pH KCl	ΔpH	pH H <sub>2</sub> O	pH KCl	ΔpH
Control	5.41 a	4.41 a	-1.00	6.39 b	5.27 bc	-1.12	5.61 b	4.08 ab	-1.53
Biochar 15 t ha <sup>-1</sup>	5.45 a	4.43 a	-1.02	6.54 b	4.28 a	-2.26	5.67 b	4.16 ab	-1.51
Biochar 30 t ha <sup>-1</sup>	5.49 a	4.44 a	-1.05	6.66 b	4.30 a	-2.36	5.71 b	3.82 a	-1.89
Compost 15 t ha <sup>-1</sup>	5.54 a	4.54 a	-1.00	6.43 b	4.67 ab	-1.76	6.43 c	4.53 c	-1.90
Compost 30 t ha <sup>-1</sup>	5.82 b	4.93 b	-0.89	6.58 b	4.98 b	-1.60	6.50 c	4.33 bc	-2.17
CaCO <sub>3</sub> 2 t ha <sup>-1</sup>	5.74 b	4.69 b	-1.05	6.95 bc	4.97 b	-1.98	6.26 c	4.81 cd	-1.45
CaCO <sub>3</sub> 4 t ha <sup>-1</sup>	5.78 b	4.79 b	-0.99	7.35 c	5.36 c	-1.99	6.35 c	5.02 d	-1.33
SP-36 2 t ha <sup>-1</sup>	5.42 a	4.36 a	-1.06	5.81 a	4.29 a	-1.52	5.15 a	4.01 a	-1.14
SP-36 4 t ha <sup>-1</sup>	5.39 a	4.32 a	-1.07	5.63 a	4.29 a	-1.34	4.92 a	3.93 a	-0.99

Values followed by the same letter in each column are not significantly different (LSD 0.05)




Figure 1. Status of pH<sub>o</sub> of three soil orders on various amendment treatments in sub-optimal dryland




Figure 2. The value of  $pH_{H_2O} - pH$  of three soil orders on various amendment treatments in sub-optimal dryland




Figure 3. The values of soil CEC of three soil orders on various amendment treatments in sub-optimal dryland




Figure 4. The values of soil ECEC of three soil orders on various amendment treatments in sub-optimal dryland